

## General

### Guideline Title

2013 ACCF/AHA guideline for the management of ST-elevation myocardial infarction. A report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines.

### Bibliographic Source(s)

O'Gara PT, Kushner FG, Ascheim DD, Casey DE Jr, Chung MK, de Lemos JA, Ettinger SM, Fang JC, Fesmire FM, Franklin BA, Granger CB, Krumholz HM, Linderbaum JA, Morrow DA, Newby LK, Ornato JP, Ou N, Radford MJ, Tamis-Holland JE, Tommaso CL, Tracy CM, Woo YJ, Zhao DX. 2013 ACCF/AHA guideline for the management of ST-elevation myocardial infarction: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. J Am Coll Cardiol. 2013 Jan 29;61(4):e78-140. [656 references] [PubMed](#)

### Guideline Status

This is the current release of the guideline.

This guideline updates previous versions: Antman EM, Anbe DT, Armstrong PW, Bates ER, Green LA, Hand M, Hochman JS, Krumholz HM, Kushner FG, Lamas GA, Mullany CJ, Ornato JP, Pearle DL, Sloan MA, Smith SC Jr. ACC/AHA guidelines for the management of patients with ST-elevation myocardial infarction. A report of the Am Coll of Cardiol/Am Heart Assoc Task Force on Practice Guidelines (Committee to revise the 1999 guidelines). Bethesda (MD): American College of Cardiology (ACC), American Heart Association (AHA); 2004. 211 p.

Canadian Cardiovascular Society, American Academy of Family Physicians, American College of Cardiology, American Heart Association, Antman EM, Hand M, Armstrong PW, Bates ER, Green LA, Halasyamani LK, Hochman JS, Krumholz HM, Lamas GA, Mullany CJ, Pearle DL, Sloan MA, Smith SC Jr, Anbe DT, Kushner FG, Ornato JP, Pearle DL, Sloan MA, Jacobs AK, Adams CD, Anderson JL, Buller CE, Creager MA, Ettinger SM, Halperin JL, Hunt SA, Lytle BW, Nishimura R, Page RL, Riegel B, Tarkington LG, Yancy CW. 2007 focused update of the ACC/AHA 2004 guidelines for the management of patients with ST-elevation myocardial infarction: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. J Am Coll Cardiol 2008 Jan 15;51(2):210-47.

## Recommendations

### Major Recommendations

Definitions for the level of the evidence (A-C) and classes of recommendations (I-III) are provided at the end of the "Major Recommendations" field.

#### Onset of Myocardial Infarction

Community Preparedness and System Goals for Reperfusion Therapy

See Figure 2, "Reperfusion therapy for patients with STEMI," in original guideline document.

#### Class I

1. All communities should create and maintain a regional system of STEMI care that includes assessment and continuous quality improvement of emergency medical services (EMS) and hospital-based activities. Performance can be facilitated by participating in programs such as Mission: Lifeline and the Door-to-Balloon (D2B) Alliance (Le May et al., 2008; Aguirre et al., 2008; Henry et al., 2007; Jollis et al., 2007). (Level of Evidence: B)
2. Performance of a 12-lead electrocardiogram (ECG) by EMS personnel at the site of first medical contact (FMC) is recommended in patients with symptoms consistent with STEMI (Sørensen et al., 2011; Le May et al., 2008; Rokos et al., 2009; Dieker et al., 2010; Diercks et al., 2009). (Level of Evidence: B)
3. Reperfusion therapy should be administered to all eligible patients with STEMI with symptom onset within the prior 12 hours (Fibrinolytic Therapy Trialists' [FTT] Collaborative Group, 1994; Keeley, Boura, & Grines, 2003). (Level of Evidence: A)
4. Primary percutaneous coronary intervention (PCI) is the recommended method of reperfusion when it can be performed in a timely fashion by experienced operators (Keeley, Boura, & Grines, 2003; Andersen et al., "Danish multicenter," 2003; Dalby et al., 2003). (Level of Evidence: A)
5. EMS transport directly to a PCI-capable hospital for primary PCI is the recommended triage strategy for patients with STEMI, with an ideal FMC-to-device time system goal of 90 minutes or less\* (Sørensen et al., 2011; Le May et al., 2008; Rokos et al., 2009). (Level of Evidence: B)
6. Immediate transfer to a PCI-capable hospital for primary PCI is the recommended triage strategy for patients with STEMI who initially arrive at or are transported to a non-PCI-capable hospital, with an FMC-to-device time system goal of 120 minutes or less\* (Andersen et al., "Danish multicenter," 2003; Dalby et al., 2003; Andersen et al., "A comparison," 2003; Nielsen et al., 2011). (Level of Evidence: B)
7. In the absence of contraindications, fibrinolytic therapy should be administered to patients with STEMI at non-PCI-capable hospitals when the anticipated FMC-to-device time at a PCI-capable hospital exceeds 120 minutes because of unavoidable delays (FTT Collaborative Group, 1994; Nallamothu & Bates, 2003; Pinto et al., 2006). (Level of Evidence: B)
8. When fibrinolytic therapy is indicated or chosen as the primary reperfusion strategy, it should be administered within 30 minutes of hospital arrival\* (Boersma et al., 1996; Chareonthaitawee et al., 2000; McNamara et al., 2007; Milavetz et al., 1998; Newby et al., 1996). (Level of Evidence: B)

\*The proposed time windows are system goals. For any individual patient, every effort should be made to provide reperfusion therapy as rapidly as possible.

#### Class IIa

1. Reperfusion therapy is reasonable for patients with STEMI and symptom onset within the prior 12 to 24 hours who have clinical and/or ECG evidence of ongoing ischemia. Primary PCI is the preferred strategy in this population (FTT Collaborative Group, 1994; Schömig et al., 2005; Gierlotka et al., 2011). (Level of Evidence: B)

#### The Relationship Between Sudden Cardiac Death and STEMI

#### *Evaluation and Management of Patients with STEMI and Out-of-Hospital Cardiac Arrest*

#### Class I

1. Therapeutic hypothermia should be started as soon as possible in comatose patients with STEMI and out-of-hospital cardiac arrest caused by ventricular fibrillation (VF) or pulseless ventricular tachycardia (VT), including patients who undergo primary PCI (Peberdy et al., 2010; Bernard, 2002; Hypothermia after Cardiac Arrest Study Group, 2002). (Level of Evidence: B)
2. Immediate angiography and PCI when indicated should be performed in resuscitated out-of-hospital cardiac arrest patients whose initial ECG shows STEMI (Nichol et al., 2010; Bendz et al., 2004; Borger van der Burg et al., 2003; Bulut et al., 2000; Garot et al., 2007; Gorjup et al., 2007; Hosmane et al., 2009; Kahn et al., 1995; Keelan et al., 2003; Kern & Rahman, 2010; Marcusohn et al., 2007; Pleskot et al., 2008; Quintero-Moran et al., 2006; Richling et al., 2007; Spaulding et al., 1997; Werling et al., 2007). (Level of Evidence: B)

#### Reperfusion at a PCI-Capable Hospital

#### Primary PCI

#### *Primary PCI in STEMI*

See Table 2 in the original guideline document for a summary of recommendations from this section.

#### Class I

1. Primary PCI should be performed in patients with STEMI and ischemic symptoms of less than 12 hours' duration (Keeley, Boura, & Grines, 2003; Zijlstra et al., 1999; "A clinical trial comparing primary coronary angioplasty," 1997). (Level of Evidence: A)
2. Primary PCI should be performed in patients with STEMI and ischemic symptoms of less than 12 hours' duration who have contraindications to fibrinolytic therapy, irrespective of the time delay from FMC (Grzybowski et al., 2003; Zahn et al., 1999). (Level of Evidence: B)
3. Primary PCI should be performed in patients with STEMI and cardiogenic shock or acute severe heart failure (HF), irrespective of time delay from myocardial infarction (MI) onset (Hochman et al., 1999; Hochman et al., "Coronary intervention," 2006; Thune et al., 2005; Wu et al., 2002). (Level of Evidence: B)

#### Class IIa

1. Primary PCI is reasonable in patients with STEMI if there is clinical and/or ECG evidence of ongoing ischemia between 12 and 24 hours after symptom onset (Schömig et al., 2005; Gierlotka et al., 2011). (Level of Evidence: B)

#### Class III: Harm

1. PCI should not be performed in a noninfarct artery at the time of primary PCI in patients with STEMI who are hemodynamically stable (Hannan et al., 2010; Toma et al., 2010; Vlaar et al., 2011). (Level of Evidence: B)

#### Aspiration Thrombectomy

#### Class IIa

1. Manual aspiration thrombectomy is reasonable for patients undergoing primary PCI (Vlaar et al., 2008; Stone et al., 2012; Bavry, Kumbhani, & Bhatt 2008; Sardella et al., 2009). (Level of Evidence: B)

#### Use of Stents in Primary PCI

#### *Use of Stents in Patients with STEMI*

#### Class I

1. Placement of a stent (bare-metal stent [BMS] or drug-eluting stent [DES]) is useful in primary PCI for patients with STEMI (Nordmann et al., 2004; Zhu et al., 2001). (Level of Evidence: A)
2. BMS† should be used in patients with high bleeding risk, inability to comply with 1 year of dual antiplatelet therapy (DAPT), or anticipated invasive or surgical procedures in the next year. (Level of Evidence: C)

†Balloon angioplasty without stent placement may be used in selected patients.

#### Class III: Harm

1. DES should not be used in primary PCI for patients with STEMI who are unable to tolerate or comply with a prolonged course of DAPT because of the increased risk of stent thrombosis with premature discontinuation of one or both agents (Spertus et al., 2006; KaÅ, uza et al., 2000; Grines et al., 2007; Park et al., 2006; Jeremias et al., 2004; Pfisterer et al., 2006; Nasser, Kapeliovich, & Markiewicz, 2005). (Level of Evidence: B)

#### Adjunctive Antithrombotic Therapy for Primary PCI

#### *Antiplatelet Therapy to Support Primary PCI for STEMI*

See Table 3 in the original guideline document for a summary of recommendations from this section.

#### Class I

1. Aspirin 162 to 325 mg should be given before primary PCI (Jolly et al., 2009; Barnathan et al., 1987; CURRENT-OASIS 7 Investigators, 2010). (Level of Evidence: B)
2. After PCI, aspirin should be continued indefinitely (Antithrombotic Trialists' Collaboration, 2002; Schömig et al., 1996; Smith et al., 2011). (Level of Evidence: A)

3. A loading dose of a P2Y<sub>12</sub>receptor inhibitor should be given as early as possible or at time of primary PCI to patients with STEMI. Options include:
  - a. Clopidogrel 600 mg (CURRENT-OASIS 7 Investigators et al., 2010; Patti et al., 2011; Mehta et al., 2010) (Level of Evidence: B)
  - b. Prasugrel 60 mg (Wiviott et al., 2007) (Level of Evidence: B)
  - c. Ticagrelor 180 mg (Steg et al., 2010) (Level of Evidence: B)
4. P2Y<sub>12</sub>inhibitor therapy should be given for 1 year to patients with STEMI who receive a stent (BMS or DES) during primary PCI using the following maintenance doses:
  - a. Clopidogrel 75 mg daily (Wiviott et al., 2007; Montalescot et al., 2009) (Level of Evidence: B)
  - b. Prasugrel 10 mg daily (Montalescot et al., 2009) (Level of Evidence: B)
  - c. Ticagrelor 90 mg twice a day<sup>‡</sup> (Steg et al., 2010) (Level of Evidence: B)

<sup>‡</sup>The recommended maintenance dose of aspirin to be used with ticagrelor is 81 mg daily.

#### Class IIa

1. It is reasonable to use 81 mg of aspirin per day in preference to higher maintenance doses after primary PCI (CURRENT-OASIS 7 Investigators, 2010; Antithrombotic Trialists' Collaboration, 2002; Serebruany et al., 2005; Steinhubl et al., 2009). (Level of Evidence: B)
2. It is reasonable to begin treatment with an intravenous glycoprotein (GP) IIb/IIIa receptor antagonist such as abciximab (Brenner et al., 1998; Stone et al., 2002; Montalescot et al., 2001) (Level of Evidence: A), high-bolus-dose tirofiban (ten Berg et al., 2010; Valgimigli et al., 2008) (Level of Evidence: B), or double-bolus eptifibatide (Akerblom et al., 2010) (Level of Evidence: B) at the time of primary PCI (with or without stenting or clopidogrel pretreatment) in selected patients with STEMI who are receiving unfractionated heparin (UFH).

#### Class IIb

1. It may be reasonable to administer intravenous GP IIb/IIIa receptor antagonist in the precatheterization laboratory setting (e.g., ambulance, emergency department) to patients with STEMI for whom primary PCI is intended (Ellis et al., 2004; ten Berg et al., 2010; Ellis et al., 2009; Montalescot et al., 2004; Maioli et al., 2007; Keeley, Boura, & Grines, 2006; Van't Hof et al., 2008; El Khoury et al., 2010; De Luca et al., 2011). (Level of Evidence: B)
2. It may be reasonable to administer intracoronary abciximab to patients with STEMI undergoing primary PCI (Stone et al., 2012; Mehilli et al., 2009; Bellandi et al., 2004; Romagnoli et al., 2005; Iversen, Galatius, & Jensen, 2008; Kakkar et al., 2004; Wöhrle et al., 2003; Bertrand et al., 2010). (Level of Evidence: B)
3. Continuation of a P2Y<sub>12</sub> inhibitor beyond 1 year may be considered in patients undergoing DES placement. (Level of Evidence: C)

#### Class III: Harm

1. Prasugrel should not be administered to patients with a history of prior stroke or transient ischemic attack (Wiviott et al., 2007). (Level of Evidence: B)

#### *Anticoagulant Therapy to Support Primary PCI*

#### Class I

1. For patients with STEMI undergoing primary PCI, the following supportive anticoagulant regimens are recommended:
  - a. UFH, with additional boluses administered as needed to maintain therapeutic activated clotting time levels, taking into account whether a GP IIb/IIIa receptor antagonist has been administered (Level of Evidence: C)
  - b. Bivalirudin with or without prior treatment with UFH (Stone et al., 2008) (Level of Evidence: B)

#### Class IIa

1. In patients with STEMI undergoing PCI who are at high risk of bleeding, it is reasonable to use bivalirudin monotherapy in preference to the combination of UFH and a GP IIb/IIIa receptor antagonist (Stone et al., 2008). (Level of Evidence: B)

#### Class III: Harm

1. Fondaparinux should not be used as the sole anticoagulant to support primary PCI because of the risk of catheter thrombosis (Yusuf et al., 2006). (Level of Evidence: B)

#### Reperfusion at a Non-PCI-Capable Hospital

## Fibrinolytic Therapy When There Is an Anticipated Delay to Performing Primary PCI within 120 Minutes of FMC

See Table 4 in the original guideline document for a summary of recommendations from this section.

### Class I

1. In the absence of contraindications, fibrinolytic therapy should be given to patients with STEMI and onset of ischemic symptoms within the previous 12 hours when it is anticipated that primary PCI cannot be performed within 120 minutes of FMC (FTT Collaborative Group, 1994; "Effect of intravenous APSAC," 1988; "Randomised trial of late thrombolysis," 1993; "Randomised trial of intravenous streptokinase," 1988; "Late Assessment of Thrombolytic Efficacy [LATE] study," 1993; Rossi & Bolognese, 1991; "A prospective trial of intravenous streptokinase," 1986). (Level of Evidence: A)

### Class IIa

1. In the absence of contraindications and when PCI is not available, fibrinolytic therapy is reasonable for patients with STEMI if there is clinical and/or electrocardiographic evidence of ongoing ischemia within 12 to 24 hours of symptom onset and a large area of myocardium at risk or hemodynamic instability. (Level of Evidence: C)

### Class III: Harm

1. Fibrinolytic therapy should not be administered to patients with ST depression except when a true posterior (inferobasal) MI is suspected or when associated with ST elevation in lead aVR (de Winter et al., 2008; Jong et al., 2006; FTT Collaborative Group, 1994; "Early effects of tissue-type plasminogen activator," 1993; Barrabés et al., 2003). (Level of Evidence: B)

## *Adjunctive Antithrombotic Therapy with Fibrinolysis*

See Table 7 in the original guideline document for a summary of recommendations from this section.

### Class I

1. Aspirin (162- to 325-mg loading dose) and clopidogrel (300-mg loading dose for patients  $\leq 75$  years of age, 75-mg dose for patients  $> 75$  years of age) should be administered to patients with STEMI who receive fibrinolytic therapy ("Randomised trial of intravenous streptokinase," 1988; Chen et al., "Addition of clopidogrel," 2005; Sabatine et al., 2005). (Level of Evidence: A)
2. Aspirin should be continued indefinitely ("Randomised trial of intravenous streptokinase," 1988; Chen et al., "Addition of clopidogrel," 2005; Sabatine et al., 2005) (Level of Evidence: A) and clopidogrel (75 mg daily) should be continued for at least 14 days (Chen et al., "Addition of clopidogrel," 2005; Sabatine et al., 2005) (Level of Evidence: A) and up to 1 year (Level of Evidence: C) in patients with STEMI who receive fibrinolytic therapy.

### Class IIa

1. It is reasonable to use aspirin 81 mg per day in preference to higher maintenance doses after fibrinolytic therapy (Antithrombotic Trialists' Collaboration, 2002; Smith et al., 2011; Serebruany et al., 2005; Steinhubl et al., 2009). (Level of Evidence: B)

## *Adjunctive Anticoagulant Therapy with Fibrinolysis*

### Class I

1. Patients with STEMI undergoing reperfusion with fibrinolytic therapy should receive anticoagulant therapy for a minimum of 48 hours, and preferably for the duration of the index hospitalization, up to 8 days or until revascularization if performed ("An international randomized trial," 1993; Antman et al., 2006). (Level of Evidence: A) Recommended regimens include:
  - a. UFH administered as a weight-adjusted intravenous bolus and infusion to obtain an activated partial thromboplastin time of 1.5 to 2.0 times control, for 48 hours or until revascularization (Level of Evidence: C)
  - b. Enoxaparin administered according to age, weight, and creatinine clearance, given as an intravenous bolus, followed in 15 minutes by subcutaneous injection for the duration of the index hospitalization, up to 8 days or until revascularization (Antman et al., 2006; Assessment of the Safety and Efficacy of a New Thrombolytic Regimen [ASSENT]-3, 2001; Ross et al., 2001; Antman et al., 2002) (Level of Evidence: A)
  - c. Fondaparinux administered with initial intravenous dose, followed in 24 hours by daily subcutaneous injections if the estimated creatinine clearance is greater than 30 mL/min, for the duration of the index hospitalization, up to 8 days or until revascularization (Yusuf et al., 2006) (Level of Evidence: B)

## Transfer to a PCI-Capable Hospital after Fibrinolytic Therapy

### *Transfer of Patients with STEMI to a PCI-Capable Hospital for Coronary Angiography after Fibrinolytic Therapy*

See Table 8 in the original guideline for a summary of recommendations from this section. See Online Data Supplement 4 for additional data on early catheterization and rescue PCI for fibrinolytic failure in the stent era; and Online Data Supplement 5 for additional data on early catheterization and PCI after fibrinolysis in the stent era (see the "Availability of Companion Documents" field).

#### Class I

1. Immediate transfer to a PCI-capable hospital for coronary angiography is recommended for suitable patients with STEMI who develop cardiogenic shock or acute severe HF, irrespective of the time delay from MI onset (Hochman et al., 2001). (Level of Evidence: B)

#### Class IIa

1. Urgent transfer to a PCI-capable hospital for coronary angiography is reasonable for patients with STEMI who demonstrate evidence of failed reperfusion or reocclusion after fibrinolytic therapy (Sutton et al., 2004; Gershlick et al., 2005; Wijeyesundera et al., 2007; Collet et al., 2006). (Level of Evidence: B)
2. Transfer to a PCI-capable hospital for coronary angiography is reasonable for patients with STEMI who have received fibrinolytic therapy even when hemodynamically stable<sup>§</sup> and with clinical evidence of successful reperfusion. Angiography can be performed as soon as logistically feasible at the receiving hospital, and ideally within 24 hours, but should not be performed within the first 2 to 3 hours after administration of fibrinolytic therapy (Böhrer et al., 2010; Borgia et al., 2010; Cantor et al., 2009; Di Mario et al., 2008; Fernandez-Avilés et al., 2004; White, 2008). (Level of Evidence: B)

<sup>§</sup>Although individual circumstances will vary, clinical stability is defined by the absence of low output, hypotension, persistent tachycardia, apparent shock, high-grade ventricular or symptomatic supraventricular tachyarrhythmias, and spontaneous recurrent ischemia.

#### Delayed Invasive Management

### Coronary Angiography in Patients Who Initially Were Managed with Fibrinolytic Therapy or Who Did Not Receive Reperfusion

See Table 9 in the original guideline document for a summary of recommendations from this section.

#### Class I

1. Cardiac catheterization and coronary angiography with intent to perform revascularization should be performed after STEMI in patients with any of the following:
  - a. Cardiogenic shock or acute severe HF that develops after initial presentation (Wu et al., 2002; Hochman et al., 2001; Steg et al., 2008; Steg et al., "Determinants and prognostic," 2004) (Level of Evidence: B)
  - b. Intermediate- or high-risk findings on predischARGE noninvasive ischemia testing (Erne et al., 2007; Madsen et al., 1997) (Level of Evidence: B)
  - c. Myocardial ischemia that is spontaneous or provoked by minimal exertion during hospitalization. (Level of Evidence: C)

#### Class IIa

1. Coronary angiography with intent to perform revascularization is reasonable for patients with evidence of failed reperfusion or reocclusion after fibrinolytic therapy. Angiography can be performed as soon as logistically feasible (Sutton et al., 2004; Gershlick et al., 2005; Wijeyesundera et al., 2007; Collet et al., 2006). (Level of Evidence: B)
2. Coronary angiography is reasonable before hospital discharge in stable<sup>§</sup> patients with STEMI after successful fibrinolytic therapy. Angiography can be performed as soon as logistically feasible, and ideally within 24 hours, but should not be performed within the first 2 to 3 hours after administration of fibrinolytic therapy (Böhrer et al., 2010; Borgia et al., 2010; Cantor et al., 2009; Di Mario et al., 2008; Fernandez-Avilés et al., 2004; White, 2008; D'Souza et al., 2011). (Level of Evidence: B)

<sup>§</sup>Although individual circumstances will vary, clinical stability is defined by the absence of low output, hypotension, persistent tachycardia, apparent shock, high-grade ventricular or symptomatic supraventricular tachyarrhythmias, and spontaneous recurrent ischemia.

### PCI of an Infarct Artery in Patients Who Initially Were Managed with Fibrinolysis or Who Did Not Receive Reperfusion Therapy

See Table 10 in the original guideline document for a summary of recommendations from this section.

#### Class I

1. PCI of an anatomically significant stenosis in the infarct artery should be performed in patients with suitable anatomy and any of the following:
  - a. Cardiogenic shock or acute severe HF (Hochman et al., 2001) (Level of Evidence: B)
  - b. Intermediate- or high-risk findings on predischARGE noninvasive ischemia testing (Erne et al., 2007; Madsen et al., 1997) (Level of Evidence: C)
  - c. Myocardial ischemia that is spontaneous or provoked by minimal exertion during hospitalization (Level of Evidence: C)

#### Class IIa

1. Delayed PCI is reasonable in patients with STEMI and evidence of failed reperfusion or reocclusion after fibrinolytic therapy. PCI can be performed as soon as logistically feasible at the receiving hospital (Gibson et al., 1999; Gibson et al., 2002; Sutton et al., 2004; Sutton et al., 2000) (Level of Evidence: B)
2. Delayed PCI of a significant stenosis in a patent infarct artery is reasonable in stable<sup>§</sup> patients with STEMI after fibrinolytic therapy. PCI can be performed as soon as logistically feasible at the receiving hospital, and ideally within 24 hours, but should not be performed within the first 2 to 3 hours after administration of fibrinolytic therapy (Böhmer et al., 2010; Borgia et al., 2010; Cantor et al., 2009; Fernandez-Avilés et al., 2004; White, 2008). (Level of Evidence: B)

<sup>§</sup>Although individual circumstances will vary, clinical stability is defined by the absence of low output, hypotension, persistent tachycardia, apparent shock, high-grade ventricular or symptomatic supraventricular tachyarrhythmias, and spontaneous recurrent ischemia.

#### Class IIb

1. Delayed PCI of a significant stenosis in a patent infarct artery greater than 24 hours after STEMI may be considered as part of an invasive strategy in stable<sup>§</sup> patients (Hochman et al., "Coronary intervention," 2006; Erne et al., 2007; Madsen et al., 1997; Gibson et al., 2003; D'Souza et al., 2011; Gupta et al., 2003; Ioannidis & Katritsis, 2007; Steg et al., "DECOPI," 2004; Wilson et al., 2001). (Level of Evidence: B)

<sup>§</sup>Although individual circumstances will vary, clinical stability is defined by the absence of low output, hypotension, persistent tachycardia, apparent shock, high-grade ventricular or symptomatic supraventricular tachyarrhythmias, and spontaneous recurrent ischemia.

#### Class III: No Benefit

1. Delayed PCI of a totally occluded infarct artery greater than 24 hours after STEMI should not be performed in asymptomatic patients with 1- or 2-vessel disease if they are hemodynamically and electrically stable and do not have evidence of severe ischemia (Hochman et al., "Coronary intervention," 2006; Ioannidis & Katritsis, 2007). (Level of Evidence: B)

#### PCI of a Noninfarct Artery before Hospital Discharge

##### Class I

1. PCI is indicated in a noninfarct artery at a time separate from primary PCI in patients who have spontaneous symptoms of myocardial ischemia. (Level of Evidence: C)

##### Class IIa

1. PCI is reasonable in a noninfarct artery at a time separate from primary PCI in patients with intermediate- or high-risk findings on noninvasive testing (Hannan et al., 2010; Erne et al., 2007; Madsen et al., 1997). (Level of Evidence: B)

#### Adjunctive Antithrombotic Therapy to Support Delayed PCI after Fibrinolytic Therapy

See Table 11 in the original guideline document for a summary of recommendations from this section.

#### *Antiplatelet Therapy to Support PCI After Fibrinolytic Therapy*

##### Class I

1. After PCI, aspirin should be continued indefinitely (CURRENT-OASIS 7 Investigators, 2010; Antithrombotic Trialists' Collaboration, 2002; Smith et al., 2011; Mehta et al., 2010; Chen et al., "Addition of clopidogrel," 2005; Sabatine et al., 2005). (Level of Evidence: A)
2. Clopidogrel should be provided as follows:
  - a. A 300-mg loading dose should be given before or at the time of PCI to patients who did not receive a previous loading dose and who are undergoing PCI within 24 hours of receiving fibrinolytic therapy (Level of Evidence: C)

- b. A 600-mg loading dose should be given before or at the time of PCI to patients who did not receive a previous loading dose and who are undergoing PCI more than 24 hours after receiving fibrinolytic therapy (Level of Evidence: C)
- c. A dose of 75 mg daily should be given after PCI (Wiviott et al., 2007; Montalescot et al., 2009; Chen et al., "Addition of clopidogrel," 2005; Sabatine et al., 2005) (Level of Evidence: C)

#### Class IIa

1. After PCI, it is reasonable to use 81 mg of aspirin per day in preference to higher maintenance doses (CURRENT-OASIS 7 Investigators, 2010; Mehta et al., 2010; Serebruany et al., 2005; Steinhubl et al., 2009). (Level of Evidence: B)
2. Prasugrel, in a 60-mg loading dose, is reasonable once the coronary anatomy is known in patients who did not receive a previous loading dose of clopidogrel at the time of administration of a fibrinolytic agent, but prasugrel should not be given sooner than 24 hours after administration of a fibrin-specific agent or 48 hours after administration of a non-fibrin-specific agent (Wiviott et al., 2007; Montalescot et al., 2009). (Level of Evidence: B)
3. Prasugrel, in a 10-mg daily maintenance dose, is reasonable after PCI (Wiviott et al., 2007; Montalescot et al., 2009). (Level of Evidence: B)

#### Class III: Harm

1. Prasugrel should not be administered to patients with a history of prior stroke or transient ischemic attack (Wiviott et al., 2007). (Level of Evidence: B)

#### *Anticoagulant Therapy to Support PCI after Fibrinolytic Therapy*

#### Class I

1. For patients with STEMI undergoing PCI after receiving fibrinolytic therapy with intravenous UFH, additional boluses of intravenous UFH should be administered as needed to support the procedure, taking into account whether GP IIb/IIIa receptor antagonists have been administered. (Level of Evidence: C)
2. For patients with STEMI undergoing PCI after receiving fibrinolytic therapy with enoxaparin, if the last subcutaneous dose was administered within the prior 8 hours, no additional enoxaparin should be given; if the last subcutaneous dose was administered between 8 and 12 hours earlier, enoxaparin 0.3 mg/kg IV should be given (Antman et al., 2002; Gibson et al., 2007). (Level of Evidence: B)

#### Class III: Harm

1. Fondaparinux should not be used as the sole anticoagulant to support PCI. An additional anticoagulant with anti-IIa activity should be administered because of the risk of catheter thrombosis (Yusuf et al., 2006). (Level of Evidence: C)

#### Coronary Artery Bypass Graft (CABG) Surgery

##### CABG in Patients with STEMI

#### Class I

1. Urgent CABG is indicated in patients with STEMI and coronary anatomy not amenable to PCI who have ongoing or recurrent ischemia, cardiogenic shock, severe HF, or other high-risk features (Caracciolo et al., 1995; Hochman et al., 2000; American College of Cardiology Foundation et al., 2011). (Level of Evidence: B)
2. CABG is recommended in patients with STEMI at time of operative repair of mechanical defects (Dalrymple-Hay et al., 1998; Menon et al., 2000; Slater et al., 2000; Tavakoli et al., 2002; Thompson et al., 2000). (Level of Evidence: B)

#### Class IIa

1. The use of mechanical circulatory support is reasonable in patients with STEMI who are hemodynamically unstable and require urgent CABG. (Level of Evidence: C)

#### Class IIb

1. Emergency CABG within 6 hours of symptom onset may be considered in patients with STEMI who do not have cardiogenic shock and are not candidates for PCI or fibrinolytic therapy. (Level of Evidence: C)

#### Timing of Urgent CABG in Patients with STEMI in Relation to Use of Antiplatelet Agents



## Class I

1. Aspirin should not be withheld before urgent CABG (Jacob et al., 2011). (Level of Evidence: C)
2. Clopidogrel or ticagrelor should be discontinued at least 24 hours before urgent on-pump CABG, if possible (Kim et al., 2008; Held et al., 2011; Nijjer et al., 2011; Barker & Anderson, 2009; Ebrahimi et al., 2009). (Level of Evidence: B)
3. Short-acting intravenous GP IIb/IIIa receptor antagonists (eptifibatide, tirofiban) should be discontinued at least 2 to 4 hours before urgent CABG (Bizzarri et al., 2001; Dyke et al., 2000). (Level of Evidence: B)
4. Abciximab should be discontinued at least 12 hours before urgent CABG (Fernandez-Avilés et al., 2004). (Level of Evidence: B)

## Class IIb

1. Urgent off-pump CABG within 24 hours of clopidogrel or ticagrelor administration might be considered, especially if the benefits of prompt revascularization outweigh the risks of bleeding (Held et al., 2011; Shim et al., 2007; Woo, Grand, & Valettas, 2003; Maltais, Perrault, & Do, 2008). (Level of Evidence: B)
2. Urgent CABG within 5 days of clopidogrel or ticagrelor administration or within 7 days of prasugrel administration might be considered, especially if the benefits of prompt revascularization outweigh the risks of bleeding. (Level of Evidence: C)

## Routine Medical Therapies

### Beta Blockers

## Class I

1. Oral beta blockers should be initiated in the first 24 hours in patients with STEMI who do not have any of the following: signs of HF, evidence of a low-output state, increased risk for cardiogenic shock<sup>†</sup>, or other contraindications to use of oral beta blockers (PR interval more than 0.24 seconds, second- or third-degree heart block, active asthma, or reactive airways disease) (Chen et al., "Early intravenous," 2005; Roberts et al., 1991; "Randomised trial of intravenous atenolol," 1986). (Level of Evidence: B)
2. Beta blockers should be continued during and after hospitalization for all patients with STEMI and with no contraindications to their use ("A randomized trial of propranolol," 1982; Freemantle et al., 1999). (Level of Evidence: B)
3. Patients with initial contraindications to the use of beta blockers in the first 24 hours after STEMI should be reevaluated to determine their subsequent eligibility. (Level of Evidence: C)

<sup>†</sup>Risk factors for cardiogenic shock (the greater the number of risk factors present, the higher the risk of developing cardiogenic shock) are age >70 years, systolic BP <120 mm Hg, sinus tachycardia >110 bpm or heart rate <60 bpm, and increased time since onset of symptoms of STEMI.

## Class IIa

1. It is reasonable to administer intravenous beta blockers at the time of presentation to patients with STEMI and no contraindications to their use who are hypertensive or have ongoing ischemia (Chen et al., "Early intravenous," 2005; Roberts et al., 1991; "Randomised trial of intravenous atenolol," 1986). (Level of Evidence: B)

### Renin-Angiotensin-Aldosterone System Inhibitors

## Class I

1. An angiotensin-converting enzyme (ACE) inhibitor should be administered within the first 24 hours to all patients with STEMI with anterior location, HF, or ejection fraction (EF) less than or equal to 0.40, unless contraindicated (Pfeffer et al., 1992; Ball, Hall, & Murray, 1994; Køber et al., 1995; Pfeffer et al., 1997). (Level of Evidence: A)
2. An angiotensin receptor blocker (ARB) should be given to patients with STEMI who have indications for but are intolerant of ACE inhibitors (Pfeffer et al., 2003; Maggioni & Fabbri, 2005) (Level of Evidence: B)
3. An aldosterone antagonist should be given to patients with STEMI and no contraindications who are already receiving an ACE inhibitor and beta blocker and who have an EF less than or equal to 0.40 and either symptomatic HF or diabetes mellitus (Pitt et al., 2003). (Level of Evidence: B)

## Class IIa

1. ACE inhibitors are reasonable for all patients with STEMI and no contraindications to their use ("Indications for ACE Inhibitors," 1998; "GISSI-3: effects of lisinopril," 1994; "ISIS-4: a randomised factorial trial," 1995). (Level of Evidence: A)

### Lipid Management

## Class I

1. High-intensity statin therapy should be initiated or continued in all patients with STEMI and no contraindications to its use (Cannon et al., 2006; Cholesterol Treatment Trialists' [CTT] Collaboration et al., 2010; Cannon et al., 2004). (Level of Evidence: B)

## Class IIa

1. It is reasonable to obtain a fasting lipid profile in patients with STEMI, preferably within 24 hours of presentation. (Level of Evidence: C)

## Complications after STEMI

### Cardiogenic Shock

#### *Treatment of Cardiogenic Shock*

## Class I

1. Emergency revascularization with either PCI or CABG is recommended in suitable patients with cardiogenic shock due to pump failure after STEMI irrespective of the time delay from MI onset (Hochman et al., 1999; Hochman et al., "Early revascularization," 2006; Babaev et al., 2005). (Level of Evidence: B)
2. In the absence of contraindications, fibrinolytic therapy should be administered to patients with STEMI and cardiogenic shock who are unsuitable candidates for either PCI or CABG (FTT Collaborative Group, 1994; Morrow et al., 2000; French et al., 2003). (Level of Evidence: B)

## Class IIa

1. The use of intra-aortic balloon pump counterpulsation can be useful for patients with cardiogenic shock after STEMI who do not quickly stabilize with pharmacological therapy (Barron et al., 2001; Chen et al., 2003; Sanborn et al., 2000; Sjaauw et al., 2009; Ohman et al., 2005). (Level of Evidence: B)

## Class IIb

1. Alternative left ventricular (LV) assist devices for circulatory support may be considered in patients with refractory cardiogenic shock. (Level of Evidence: C)

### Electrical Complications during the Hospital Phase of STEMI

#### *Implantable Cardioverter-Defibrillator (ICD) Therapy before Discharge*

## Class I

1. ICD therapy is indicated before discharge in patients who develop sustained VT/VF more than 48 hours after STEMI, provided the arrhythmia is not due to transient or reversible ischemia, reinfarction, or metabolic abnormalities (Wever et al., 1995; Siebels & Kuck, 1994; Connolly et al., 2000). (Level of Evidence: B)

#### *Bradycardia, Atrioventricular (AV) Block, and Intraventricular Conduction Defects*

## Pacing in STEMI

## Class I

1. Temporary pacing is indicated for symptomatic bradyarrhythmias unresponsive to medical treatment. (Level of Evidence: C)

### Pericarditis

#### *Management of Pericarditis after STEMI*

## Class I

1. Aspirin is recommended for treatment of pericarditis after STEMI (Berman, Haffajee, & Alpert, 1981). (Level of Evidence: B)

## Class IIb

1. Administration of acetaminophen, colchicine, or narcotic analgesics may be reasonable if aspirin, even in higher doses, is not effective. (Level of Evidence: C)

### Class III: Harm

1. Glucocorticoids and nonsteroidal anti-inflammatory drugs are potentially harmful for treatment of pericarditis after STEMI (Bulkley & Roberts, 1974; Silverman & Pfeifer, 1987). (Level of Evidence: B)

### Thromboembolic and Bleeding Complications

#### *Thromboembolic Complications*

### Anticoagulation

Note: These recommendations apply to patients who receive intracoronary stents during PCI for STEMI. Among individuals with STEMI who do not receive an intracoronary stent, the duration of DAPT beyond 14 days has not been studied adequately for patients who undergo balloon angioplasty alone, are treated with fibrinolysis alone, or do not receive reperfusion therapy. In this subset of patients with STEMI who do not receive an intracoronary stent, the threshold for initiation of oral anticoagulation for secondary prevention, either alone or in combination with aspirin, may be lower, especially if a shorter duration (i.e., 14 days) of DAPT is planned (Andreotti et al., 2006).

### Class I

1. Anticoagulant therapy with a vitamin K antagonist should be provided to patients with STEMI and atrial fibrillation with CHADS<sub>2</sub> (Congestive heart failure, Hypertension, Age  $\geq 75$  years, Diabetes mellitus, previous Stroke/transient ischemic attack [doubled risk weight]) score greater than or equal to 2, mechanical heart valves, venous thromboembolism, or hypercoagulable disorder. (Level of Evidence: C)
2. The duration of triple antithrombotic therapy with a vitamin K antagonist, aspirin, and a P2Y<sub>12</sub> receptor inhibitor should be minimized to the extent possible to limit the risk of bleeding.\*\* (Level of Evidence: C)

\*\*Individual circumstances will vary and depend on the indications for triple therapy and the type of stent placed during PCI. After this initial treatment period, consider therapy with a vitamin K antagonist plus a single antiplatelet agent. For patients treated with fibrinolysis, consider triple therapy for 14 days, followed by a vitamin K antagonist plus a single antiplatelet agent (You et al., 2012; Vandvik et al., 2012; Lip et al., 2010; Faxon et al., 2011).

### Class IIa

1. Anticoagulant therapy with a vitamin K antagonist is reasonable for patients with STEMI and asymptomatic LV mural thrombi. (Level of Evidence: C)

### Class IIb

1. Anticoagulant therapy may be considered for patients with STEMI and anterior apical akinesis or dyskinesis. (Level of Evidence: C)
2. Targeting vitamin K antagonist therapy to a lower international normalized ratio (e.g., 2.0 to 2.5) might be considered in patients with STEMI who are receiving DAPT. (Level of Evidence: C)

### Risk Assessment after STEMI

#### Use of Noninvasive Testing for Ischemia before Discharge

### Class I

1. Noninvasive testing for ischemia should be performed before discharge to assess the presence and extent of inducible ischemia in patients with STEMI who have not had coronary angiography and do not have high-risk clinical features for which coronary angiography would be warranted (Th  roux et al., 1979; Villella et al., 1995; Leppo et al., 1984). (Level of Evidence: B)

### Class IIb

1. Noninvasive testing for ischemia might be considered before discharge to evaluate the functional significance of a noninfarct artery stenosis previously identified at angiography. (Level of Evidence: C)
2. Noninvasive testing for ischemia might be considered before discharge to guide the postdischarge exercise prescription. (Level of Evidence: C)

### Assessment of LV Function

## Class I

1. LV ejection fraction (LVEF) should be measured in all patients with STEMI. (Level of Evidence: C)

## Assessment of Risk for Sudden Cardiac Death

## Class I

1. Patients with an initially reduced LVEF who are possible candidates for implantable cardioverter-defibrillator therapy should undergo reevaluation of LVEF 40 or more days after discharge (Epstein et al., 2008; Hohnloser et al., 2004; Steinbeck et al., 2009; Moss et al., 2002). (Level of Evidence: B)

## Post-hospitalization Plan of Care

## Class I

1. Posthospital systems of care designed to prevent hospital readmissions should be used to facilitate the transition to effective, coordinated outpatient care for all patients with STEMI (Naylor et al., 1994; Coleman et al., 2006; Young et al., 2003; Jack et al., 2009; Lappé et al., 2004). (Level of Evidence: B)
2. Exercise-based cardiac rehabilitation/secondary prevention programs are recommended for patients with STEMI (Leon et al., 2005; Suaya et al., 2009; Taylor et al., 2004; Goel et al., 2011). (Level of Evidence: B)
3. A clear, detailed, and evidence-based plan of care that promotes medication adherence, timely follow-up with the healthcare team, appropriate dietary and physical activities, and compliance with interventions for secondary prevention should be provided to patients with STEMI. (Level of Evidence: C)
4. Encouragement and advice to stop smoking and to avoid second-hand smoke should be provided to patients with STEMI (Wilson et al., 2000; Thomson & Rigotti, 2003; Dawood et al., 2008; Shah et al., 2010). (Level of Evidence: A)

## Definitions:

## Applying Classification of Recommendations and Level of Evidence

		Size of Treatment Effect					
		CLASS I  <i>Benefit &gt;&gt;&gt; Risk</i>  Procedure/Treatment SHOULD be performed/ administered	CLASS IIa  <i>Benefit &gt;&gt; Risk</i> <i>Additional studies with focused objectives needed</i>  IT IS REASONABLE to perform procedure/administer treatment	CLASS IIb  <i>Benefit ≥ Risk</i> <i>Additional studies with broad objectives needed; additional registry data would be helpful</i>  Procedure/Treatment MAY BE CONSIDERED	CLASS III <i>No Benefit</i> or Class III <i>Harm</i>		
						Procedure/Test	Treatment
					COR III: No Benefit	Not helpful	No proven benefit
					COR III: Harm	Excess Cost without Benefit or Harmful	Harmful to Patients
Estimate of Certainty (Precision) of Treatment Effect	LEVEL A  Multiple populations evaluated*  Data derived from multiple randomized clinical trials or meta-analyses	<ul style="list-style-type: none"><li>Recommendation that procedure or treatment is useful/effective</li><li>Sufficient evidence from multiple randomized trials or meta-analyses</li></ul>	<ul style="list-style-type: none"><li>Recommendation in favor of treatment or procedure being useful/effective</li><li>Some conflicting evidence from multiple randomized trials or meta-analyses</li></ul>	<ul style="list-style-type: none"><li>Recommendation's usefulness/efficacy less well established</li><li>Greater conflicting evidence from multiple randomized trials or meta-analyses</li></ul>	<ul style="list-style-type: none"><li>Recommendation that procedure or treatment is not useful/effective and may be harmful</li><li>Sufficient evidence from multiple randomized trials or meta-analyses</li></ul>		
	LEVEL B  Limited populations evaluated*  Data derived from a single randomized clinical trial or nonrandomized studies	<ul style="list-style-type: none"><li>Recommendation that procedure or treatment is useful/effective</li><li>Evidence from single randomized trial or nonrandomized studies</li></ul>	<ul style="list-style-type: none"><li>Recommendation in favor of treatment or procedure being useful/effective</li><li>Some conflicting evidence from single randomized trial or nonrandomized studies</li></ul>	<ul style="list-style-type: none"><li>Recommendation's usefulness/efficacy less well established</li><li>Greater conflicting evidence from single randomized trial or nonrandomized studies</li></ul>	<ul style="list-style-type: none"><li>Recommendation that procedure or treatment is not useful/effective and may be harmful</li><li>Evidence from single randomized trial or nonrandomized studies</li></ul>		
	LEVEL C  Very limited populations evaluated*  Only consensus opinion of experts, case studies or standard of care	<ul style="list-style-type: none"><li>Recommendation that procedure or treatment is useful/effective</li><li>Only expert opinion, case studies, or standard of care</li></ul>	<ul style="list-style-type: none"><li>Recommendation in favor of treatment or procedure being useful/effective</li><li>Only diverging expert opinion, case studies, or standard of care</li></ul>	<ul style="list-style-type: none"><li>Recommendation's usefulness/efficacy less well established</li><li>Only diverging expert opinion, case studies, or standard of care</li></ul>	<ul style="list-style-type: none"><li>Recommendation that procedure or treatment is not useful/effective and may be harmful</li><li>Only expert opinion, case studies, or standard of care</li></ul>		

A recommendation with Level of Evidence B or C does not imply that the recommendation is weak. Many important clinical questions addressed in the guidelines do not lend themselves to clinical trials. Although randomized trials are unavailable, there may be a very clear clinical consensus that a particular test or therapy is useful or effective.

\*Data available from clinical trials or registries about the usefulness/efficacy in different subpopulations, such as gender, age, history of diabetes, history of prior myocardial infarction, history of heart failure, and prior aspirin use.

## Clinical Algorithm(s)

A clinical algorithm for reperfusion therapy for patients with ST-elevation myocardial infarction (STEMI) is provided in the original guideline document.

## Scope

### Disease/Condition(s)

ST-elevation myocardial infarction (STEMI)

### Guideline Category

Diagnosis

Evaluation

Management

Risk Assessment

Treatment

### Clinical Specialty

Cardiology

Critical Care

Emergency Medicine

Family Practice

Internal Medicine

Radiology

Thoracic Surgery

### Intended Users

Advanced Practice Nurses

Emergency Medical Technicians/Paramedics

Health Care Providers

Nurses

Pharmacists

Physician Assistants

Physicians

## Guideline Objective(s)

- To review the management of patients with ST-elevation myocardial infarction (STEMI), with particular emphasis on advances in reperfusion therapy, organization of regional systems of care, transfer algorithms, evidence-based antithrombotic and medical therapies, and secondary prevention strategies to optimize patient-centered care
- To assist healthcare providers in clinical decision making by describing a range of generally acceptable approaches for the diagnosis, management, and prevention of STEMI

## Target Population

- Adults with ST-elevation myocardial infarction (STEMI)
- Adults at risk of STEMI

## Interventions and Practices Considered

### Diagnosis/Evaluation/Risk Assessment

1. Performance of a 12-lead electrocardiogram (ECG) by emergency medical services (EMS) personnel at the site of first medical contact (FMC)
2. Evaluation of patients with ST-elevation myocardial infarction (STEMI) and out-of-hospital cardiac arrest (immediate angiography)
3. Risk assessment after STEMI
  - Noninvasive testing for ischemia in patients who have not undergone angiography or who have a significant stenosis in a noninfarct artery
  - Assessment of left ventricular function
  - Assessment of risk for sudden cardiac death

### Management/Treatment

1. Creating and maintaining a regional system of STEMI care
2. Administration of reperfusion therapy to all eligible patients (preferably within 12 hours of symptom onset)
3. Primary percutaneous coronary intervention (PCI) as method of reperfusion
4. EMS—rather than private—transport to PCI-capable facility
5. Management of patients with STEMI and out-of-hospital cardiac arrest (use of therapeutic hypothermia and PCI as indicated)
6. Manual aspiration thrombectomy
7. Placement of a stent (bare-metal stent or drug-eluting stent)
8. Antiplatelet therapy to support primary PCI (aspirin, P2Y<sub>12</sub> receptor inhibitor, glycoprotein [GP] IIb/IIIa receptor antagonist)
9. Anticoagulant therapy to support primary PCI (unfractionated heparin, bivalirudin, fondaparinux) (fondaparinux not recommended as sole anticoagulant)
10. Fibrinolytic therapy if anticipated delay to performing primary PCI exceeds 120 minutes from time of first medical contact
11. Adjunctive antiplatelet or anticoagulant therapy after fibrinolytic therapy
12. Immediate transfer to a PCI-capable hospital for coronary angiography (CA) after fibrinolytic therapy whether successful or unsuccessful
13. PCI of an infarct artery with a significant stenosis in patients who initially were managed with fibrinolysis or who did not receive reperfusion therapy
14. PCI of a significant stenosis in a noninfarct artery at a time removed from primary PCI and before hospital discharge in patients without cardiogenic shock or severe heart failure
15. PCI of a noninfarct artery at time of primary PCI for patients with cardiogenic shock or severe heart failure
16. Urgent coronary artery bypass graft (CABG) when indicated
17. Timing of urgent CABG in relation to use of antiplatelet agents
18. Routine medical therapies
  - Beta-blockers
  - Angiotensin-converting enzyme (ACE) inhibitors
  - Angiotensin receptor blockers
  - Aldosterone antagonists
  - Lipid management, including statin therapy and fasting lipid profile

## 19. Management of complications after STEMI

- Treatment of cardiogenic shock (emergency revascularization with either PCI or CA, alternative left ventricular [LV] assist devices)
- Electrical complications (implantable cardioverter-defibrillator therapy, temporary pacing)
- Management of pericarditis (aspirin and analgesics)
- Thromboembolic and bleeding complications (anticoagulant therapy with a vitamin K antagonist)

## 20. Posthospital systems of care designed to prevent hospital readmissions, including detailed plans of care

## 21. Exercise-based cardiac rehabilitation/secondary prevention programs

## 22. Encouragement and advice to stop smoking and avoidance of second-hand smoke

Note: Glucocorticoids and nonsteroidal anti-inflammatory drugs not recommended for management of pericarditis.

## Major Outcomes Considered

- Morbidity and mortality due to ST-elevation myocardial infarction (STEMI)
- Primary prevention of STEMI
- Secondary prevention of cardiovascular events, including second myocardial infarction, sudden cardiac death, recurrent myocardial ischemia, stroke
- Time to treatment
- Incidence of serious bleeding or stroke
- Rates of adverse events and/or side effects of treatment options

## Methodology

### Methods Used to Collect/Select the Evidence

Hand-searches of Published Literature (Primary Sources)

Hand-searches of Published Literature (Secondary Sources)

Searches of Electronic Databases

### Description of Methods Used to Collect/Select the Evidence

The recommendations listed in this document are, whenever possible, evidence based. The current document constitutes a full revision and includes an extensive evidence review, which was conducted through November 2010, with additional selected references added through August 2012. PubMed and the Cochrane Collaboration Database were used. Searches were limited to studies conducted in human subjects and reviews and other evidence pertaining to human subjects; all were published in English. Key search words included but were not limited to: *acute coronary syndromes, percutaneous coronary intervention, coronary artery bypass graft, myocardial infarction, ST-elevation myocardial infarction, coronary stent, revascularization, anticoagulant therapy, antiplatelet therapy, antithrombotic therapy, glycoprotein IIb/IIIa inhibitor therapy, pharmacotherapy, proton-pump inhibitor, implantable cardioverter-defibrillator therapy, cardiogenic shock, fibrinolytic therapy, thrombolytic therapy, nitrates, mechanical complications, arrhythmia, angina, chronic stable angina, diabetes, chronic kidney disease, mortality, morbidity, elderly, ethics, and contrast nephropathy*. Additional searches cross-referenced these topics with the following subtopics: *percutaneous coronary intervention, coronary artery bypass graft, cardiac rehabilitation, and secondary prevention*. Additionally, the committee reviewed documents related to the subject matter previously published by the American College of Cardiology Foundation (ACCF) and American Heart Association (AHA). References selected and published in this document are representative and not all-inclusive.

### Number of Source Documents

Not stated

# Methods Used to Assess the Quality and Strength of the Evidence

Weighting According to a Rating Scheme (Scheme Given)

## Rating Scheme for the Strength of the Evidence

Applying Classification of Recommendations and Level of Evidence

		Size of Treatment Effect					
		CLASS I  <i>Benefit &gt;&gt;&gt; Risk</i>  Procedure/Treatment SHOULD be performed/ administered	CLASS IIa  <i>Benefit &gt;&gt; Risk</i> <i>Additional studies with focused objectives needed</i>  IT IS REASONABLE to perform procedure/administer treatment	CLASS IIb  <i>Benefit ≥ Risk</i> <i>Additional studies with broad objectives needed; additional registry data would be helpful</i>  Procedure/Treatment MAY BE CONSIDERED	CLASS III <i>No Benefit</i> or Class III <i>Harm</i>		
						Procedure/Test	Treatment
					COR III: No Benefit	Not helpful	No proven benefit
					COR III: Harm	Excess Cost without Benefit or Harmful	Harmful to Patients
Estimate of Certainty (Precision) of Treatment Effect	LEVEL A  Multiple populations evaluated*  Data derived from multiple randomized clinical trials or meta-analyses	<ul style="list-style-type: none"><li>Recommendation that procedure or treatment is useful/effective</li><li>Sufficient evidence from multiple randomized trials or meta-analyses</li></ul>	<ul style="list-style-type: none"><li>Recommendation in favor of treatment or procedure being useful/effective</li><li>Some conflicting evidence from multiple randomized trials or meta-analyses</li></ul>	<ul style="list-style-type: none"><li>Recommendation's usefulness/efficacy less well established</li><li>Greater conflicting evidence from multiple randomized trials or meta- analyses</li></ul>	<ul style="list-style-type: none"><li>Recommendation that procedure or treatment is not useful/effective and may be harmful</li><li>Sufficient evidence from multiple randomized trials or meta-analyses</li></ul>		
	LEVEL B  Limited populations evaluated*  Data derived from a single randomized clinical trials or nonrandomized studies	<ul style="list-style-type: none"><li>Recommendation that procedure or treatment is useful/effective</li><li>Evidence from single randomized trial or nonrandomized studies</li></ul>	<ul style="list-style-type: none"><li>Recommendation in favor of treatment or procedure being useful/effective</li><li>Some conflicting evidence from single randomized trial or nonrandomized studies</li></ul>	<ul style="list-style-type: none"><li>Recommendation's usefulness/efficacy less well established</li><li>Greater conflicting evidence from single randomized trial or nonrandomized studies</li></ul>	<ul style="list-style-type: none"><li>Recommendation that procedure or treatment is not useful/effective and may be harmful</li><li>Evidence from single randomized trial or nonrandomized studies</li></ul>		
	LEVEL C  Very limited populations evaluated*  Only consensus opinion of experts, case studies or standard of care	<ul style="list-style-type: none"><li>Recommendation that procedure or treatment is useful/effective</li><li>Only expert opinion, case studies, or standard of care</li></ul>	<ul style="list-style-type: none"><li>Recommendation in favor of treatment or procedure being useful/effective</li><li>Only diverging expert opinion, case studies, or standard of care</li></ul>	<ul style="list-style-type: none"><li>Recommendation's usefulness/efficacy less well established</li><li>Only diverging expert opinion, case studies, or standard of care</li></ul>	<ul style="list-style-type: none"><li>Recommendation that procedure or treatment is not useful/effective and may be harmful</li><li>Only expert opinion, case studies, or standard of care</li></ul>		

A recommendation with Level of Evidence B or C does not imply that the recommendation is weak. Many important clinical questions addressed in the guidelines do not lend themselves to clinical trials. Although randomized trials are unavailable, there may be a very clear clinical consensus that a particular test or therapy is useful or effective.

\*Data available from clinical trials or registries about the usefulness/efficacy in different subpopulations, such as gender, age, history of diabetes, history of prior myocardial infarction, history of heart failure, and prior aspirin use.

## Methods Used to Analyze the Evidence

Review of Published Meta-Analyses

Systematic Review with Evidence Tables

## Description of the Methods Used to Analyze the Evidence

To provide clinicians with a comprehensive set of data, whenever deemed appropriate or when published, the absolute risk difference and number needed to treat or harm are provided in the guideline, along with confidence intervals (CI) and data related to the relative treatment effects such as odds ratio (OR), relative risk (RR), hazard ratio (HR), or incidence rate ratio

## Methods Used to Formulate the Recommendations

Expert Consensus



## Description of Methods Used to Formulate the Recommendations

Experts in the subject under consideration are selected by the American College of Cardiology Foundation (ACCF) and American Heart Association (AHA) to examine subject-specific data and write guidelines in partnership with representatives from other medical organizations and specialty groups. Writing committees are asked to perform a formal literature review; weigh the strength of evidence for or against particular tests, treatments, or procedures; and include estimates of expected outcomes where such data exist. Patient-specific modifiers, comorbidities, and issues of patient preference that may influence the choice of tests or therapies are considered. When available, information from studies on cost is considered, but data on efficacy and outcomes constitute the primary basis for the recommendations contained herein.

In analyzing the data and developing recommendations and supporting text, the writing committee uses evidence-based methodologies developed by the Task Force. The Class of Recommendation (COR) is an estimate of the size of the treatment effect considering risks versus benefits in addition to evidence and/or agreement that a given treatment or procedure is or is not useful/effective or in some situations may cause harm. The Level of Evidence (LOE) is an estimate of the certainty or precision of the treatment effect. The writing committee reviews and ranks evidence supporting each recommendation with the weight of evidence ranked as LOE A, B, or C according to specific definitions that are included in the "Rating Scheme for the Strength of the Evidence" field. Studies are identified as observational, retrospective, prospective, or randomized where appropriate. For certain conditions for which inadequate data are available, recommendations are based on expert consensus and clinical experience and are ranked as LOE C. When recommendations at LOE C are supported by historical clinical data, appropriate references (including clinical reviews) are cited if available. For issues for which sparse data are available, a survey of current practice among the members of the writing committee is the basis for LOE C recommendations and no references are cited.

A new addition to this methodology is separation of the Class III recommendations to delineate whether the recommendation is determined to be of "no benefit" or is associated with "harm" to the patient. In addition, in view of the increasing number of comparative effectiveness studies, comparator verbs and suggested phrases for writing recommendations for the comparative effectiveness of one treatment or strategy versus another are included for COR I and IIa, LOE A or B only.

The writing committee was composed of experts representing cardiovascular medicine, interventional cardiology, electrophysiology, heart failure (HF), cardiac surgery, emergency medicine, internal medicine, cardiac rehabilitation, nursing, and pharmacy. The American College of Physicians, American College of Emergency Physicians, and Society for Cardiovascular Angiography and Interventions assigned official representatives.

## Rating Scheme for the Strength of the Recommendations

See the "Rating Scheme for the Strength of the Evidence" field, above.

## Cost Analysis

The guideline developers reviewed published cost analyses.

## Method of Guideline Validation

External Peer Review

Internal Peer Review

## Description of Method of Guideline Validation

This document was reviewed by 2 outside reviewers each nominated by the American College of Cardiology Foundation (ACCF) and the American Heart Association (AHA), as well as 2 reviewers each from the American College of Emergency Physicians (ACEP) and Society for Cardiovascular Angiography and Interventions (SCAI) and 22 individual content reviewers (including members from the ACCF Interventional Scientific Council and ACCF Surgeons' Scientific Council). All reviewer relationship with industry (RWI) information was distributed to the writing committee and is published in the original guideline Appendix.

This document was approved for publication by the governing bodies of the ACCF and the AHA and was endorsed by the ACEP and SCAI.

# Evidence Supporting the Recommendations

## References Supporting the Recommendations

A clinical trial comparing primary coronary angioplasty with tissue plasminogen activator for acute myocardial infarction. The Global Use of Strategies to Open Occluded Coronary Arteries in Acute Coronary Syndromes (GUSTO IIb) Angioplasty Substudy Invest. N Engl J Med. 1997 Jun 5;336(23):1621-8. [PubMed](#)

A prospective trial of intravenous streptokinase in acute myocardial infarction (I.S.A.M.). Mortality, morbidity, and infarct size at 21 days. The I.S.A.M. Study Group. N Engl J Med. 1986 Jun 5;314(23):1465-71. [PubMed](#)

A randomized trial of propranolol in patients with acute myocardial infarction. I. Mortality results. JAMA. 1982 Mar 26;247(12):1707-14. [PubMed](#)

Aguirre FV, Varghese JJ, Kelley MP, Lam W, Lucore CL, Gill JB, Page L, Turner L, Davis C, Mikell FL, Stat Heart Investigators. Rural interhospital transfer of ST-elevation myocardial infarction patients for percutaneous coronary revascularization: the Stat Heart Program. Circulation. 2008 Mar 4;117(9):1145-52. [PubMed](#)

Akerblom A, James SK, Koutouzis M, Lagerqvist B, Stenestrand U, Svennblad B, Oldgren J. Eptifibatide is noninferior to abciximab in primary percutaneous coronary intervention: results from the SCAAR (Swedish Coronary Angiography and Angioplasty Registry). J Am Coll Cardiol. 2010 Aug 3;56(6):470-5. [PubMed](#)

American College of Cardiology Foundation, American Heart Association Task Force on Practice Guidelines, American Association for Thoracic Surgery, Society of Cardiovascular Anesthesiologists, Society of Thoracic Surgeons, Hillis LD, Smith PK, Anderson JL, Bittl JA, Bridges CR, Byrne JG, Cigarroa JE, DiSesa VJ, Hiratzka LF, Hutter AM Jr, Jessen ME, Keeley EC, Lahey SJ, Lange RA, London MJ, Mack MJ, Patel MR, Puskas JD, Sabik JF, Selnes O, Shahian DM, Trost JC, Winniford MD. 2011 ACCF/AHA guideline for coronary artery bypass graft surgery: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines [trunc]. J Am Coll Cardiol. 2011 Dec 6;58(24):e123-210. [1264 references] [PubMed](#)

An international randomized trial comparing four thrombolytic strategies for acute myocardial infarction. The GUSTO investigators. N Engl J Med. 1993 Sep 2;329(10):673-82. [PubMed](#)

Andersen HR, Nielsen TT, Rasmussen K, Thuesen L, Kelbaek H, Thayssen P, Abildgaard U, Pedersen F, Madsen JK, Grande P, Villadsen AB, Krusell LR, Haghfelt T, Lomholt P, Husted SE, Vigholt E, Kjaergard HK, Mortensen LS. A comparison of coronary angioplasty with fibrinolytic therapy in acute myocardial infarction. N Engl J Med. 2003 Aug 21;349(8):733-42. [PubMed](#)

Andersen HR, Nielsen TT, Vesterlund T, Grande P, Abildgaard U, Thayssen P, Pedersen F, Mortensen LS. Danish multicenter randomized study on fibrinolytic therapy versus acute coronary angioplasty in acute myocardial infarction: rationale and design of the DANish trial in Acute Myocardial Infarction-2 (DANAMI-2). Am Heart J. 2003 Aug;146(2):234-41. [PubMed](#)

Andreotti F, Testa L, Biondi-Zoccai GG, Crea F. Aspirin plus warfarin compared to aspirin alone after acute coronary syndromes: an updated and comprehensive meta-analysis of 25,307 patients. Eur Heart J. 2006 Mar;27(5):519-26. [31 references] [PubMed](#)

Antithrombotic Trialists' Collaboration. Collaborative meta-analysis of randomised trials of antiplatelet therapy for prevention of death, myocardial infarction, and stroke in high risk patients. BMJ. 2002 Jan 12;324(7329):71-86. [PubMed](#)

Antman EM, Louwerenburg HW, Baars HF, Wesdorp JC, Hamer B, Bassand JP, Bigonzi F, Pisapia G, Gibson CM, Heidbuchel H, Braunwald E, Van de Werf F. Enoxaparin as adjunctive antithrombin therapy for ST-elevation myocardial infarction [trunc] [Erratum in Circulation 2002 Jun 11;105(23):2799]. Circulation. 2002 Apr 9;105(14):1642-9. [PubMed](#)

Antman EM, Morrow DA, McCabe CH, Murphy SA, Ruda M, Sadowski Z, Budaj A, Lopez-Sendon JL, Guneri S, Jiang F, White HD, Fox KA, Braunwald E, ExTRACT-TIMI 25 Investigators. Enoxaparin versus unfractionated heparin with fibrinolysis for ST-elevation myocardial infarction. *N Engl J Med*. 2006 Apr 6;354(14):1477-88. [PubMed](#)

Assessment of the Safety and Efficacy of a New Thrombolytic Regimen (ASSENT)-3 Investigators. Efficacy and safety of tenecteplase in combination with enoxaparin, abciximab, or unfractionated heparin: the ASSENT-3 randomised trial in acute myocardial infarction. *Lancet*. 2001 Aug 25;358(9282):605-13. [PubMed](#)

Babaev A, Frederick PD, Pasta DJ, Every N, Sichrovsky T, Hochman JS, NRMI Investigators. Trends in management and outcomes of patients with acute myocardial infarction complicated by cardiogenic shock. *JAMA*. 2005 Jul 27;294(4):448-54. [PubMed](#)

Ball SG, Hall AS, Murray GD. ACE inhibition, atherosclerosis and myocardial infarction--the AIRE Study in practice. Acute Infarction Ramipril Efficacy Study. *Eur Heart J*. 1994 May;15(Suppl B):20-5; discussion 26-30. [PubMed](#)

Barker CM, Anderson HV. Acute coronary syndromes: don't bypass the clopidogrel. *J Am Coll Cardiol*. 2009 May 26;53(21):1973-4. [PubMed](#)

Barnathan ES, Schwartz JS, Taylor L, Laskey WK, Kleaveland JP, Kussmaul WG, Hirshfeld JW Jr. Aspirin and dipyridamole in the prevention of acute coronary thrombosis complicating coronary angioplasty. *Circulation*. 1987 Jul;76(1):125-34. [PubMed](#)

Barrabes JA, Figueras J, Moure C, Cortadellas J, Soler-Soler J. Prognostic value of lead aVR in patients with a first non-ST-segment elevation acute myocardial infarction. *Circulation*. 2003 Aug 19;108(7):814-9. [PubMed](#)

Barron HV, Every NR, Parsons LS, Angeja B, Goldberg RJ, Gore JM, Chou TM, Investigators in the National Registry of Myocardial Infarction 2. The use of intra-aortic balloon counterpulsation in patients with cardiogenic shock complicating acute myocardial infarction: data from the National Registry of Myocardial Infarction 2. *Am Heart J*. 2001 Jun;141(6):933-9. [PubMed](#)

Bavry AA, Kumbhani DJ, Bhatt DL. Role of adjunctive thrombectomy and embolic protection devices in acute myocardial infarction: a comprehensive meta-analysis of randomized trials. *Eur Heart J*. 2008 Dec;29(24):2989-3001. [50 references] [PubMed](#)

Bellandi F, Maioli M, Gallopin M, Toso A, Dabizzi RP. Increase of myocardial salvage and left ventricular function recovery with intracoronary abciximab downstream of the coronary occlusion in patients with acute myocardial infarction treated with primary coronary intervention. *Catheter Cardiovasc Interv*. 2004 Jun;62(2):186-92. [PubMed](#)

Bendz B, Eritsland J, Nakstad AR, Brekke M, Klow NE, Steen PA, Mangschau A. Long-term prognosis after out-of-hospital cardiac arrest and primary percutaneous coronary intervention. *Resuscitation*. 2004 Oct;63(1):49-53. [PubMed](#)

Berman J, Haffajee CI, Alpert JS. Therapy of symptomatic pericarditis after myocardial infarction: retrospective and prospective studies of aspirin, indomethacin, prednisone, and spontaneous resolution. *Am Heart J*. 1981 Jun;101(6):750-3. [PubMed](#)

Bernard SA, Gray TW, Buist MD, Jones BM, Silvester W, Gutteridge G, Smith K. Treatment of comatose survivors of out-of-hospital cardiac arrest with induced hypothermia. *N Engl J Med*. 2002 Feb 21;346(8):557-63. [PubMed](#)

Bertrand OF, Rodes-Cabau J, Larose E, Rinfret S, Gaudreault V, Proulx G, Barbeau G, Dery JP, Gleeton O, Manh-Nguyen C, Noel B, Roy L, Costerousse O, De Larochelliere R, EARly Discharge after Transradial Stenting of Coronary Arteries in Acute. Intracoronary compared to intravenous Abciximab and high-dose bolus compared to standard dose in patients with ST-segment elevation myocardial infarction undergoing transradial primary percutaneous coronary intervention: a two-by-two factorial [trunc]. *Am J Cardiol*. 2010 Jun 1;105(11):1520-7. [PubMed](#)

Bizzarri F, Scolletta S, Tucci E, Lucidi M, Davoli G, Toscano T, Neri E, Muzzi L, Frati G. Perioperative use of tirofiban hydrochloride

(Aggrastat) does not increase surgical bleeding after emergency or urgent coronary artery bypass grafting. *J Thorac Cardiovasc Surg*. 2001 Dec;122(6):1181-5. [PubMed](#)

Boersma E, Maas AC, Deckers JW, Simoons ML. Early thrombolytic treatment in acute myocardial infarction: reappraisal of the golden hour. *Lancet*. 1996 Sep 21;348(9030):771-5. [PubMed](#)

Bohmer E, Hoffmann P, Abdelnoor M, Arnesen H, Halvorsen S. Efficacy and safety of immediate angioplasty versus ischemia-guided management after thrombolysis in acute myocardial infarction in areas with very long transfer distances results of the NORDISTEMI [trunc]. *J Am Coll Cardiol*. 2010 Jan 12;55(2):102-10. [PubMed](#)

Borger van der Burg AE, Bax JJ, Boersma E, Bootsma M, van Erven L, van der Wall EE, Schalij MJ. Impact of percutaneous coronary intervention or coronary artery bypass grafting on outcome after nonfatal cardiac arrest outside the hospital. *Am J Cardiol*. 2003 Apr 1;91(7):785-9. [PubMed](#)

Borgia F, Goodman SG, Halvorsen S, Cantor WJ, Piscione F, Le May MR, Fernandez-Aviles F, Sanchez PL, Dimopoulos K, Scheller B, Armstrong PW, Di Mario C. Early routine percutaneous coronary intervention after fibrinolysis vs. standard therapy in ST-segment elevation myocardial infarction: a meta-analysis. *Eur Heart J*. 2010 Sep;31(17):2156-69. [PubMed](#)

Brener SJ, Barr LA, Burchenal JE, Katz S, George BS, Jones AA, Cohen ED, Gaaney PC, White HJ, Cheek HB, Moses JW, Moliterno DJ, Effron MB, Topol EJ. Randomized, placebo-controlled trial of platelet glycoprotein IIb/IIIa blockade with primary angioplasty for acute myocardial infarction. ReoPro and Primary PTCA Organization and Randomized Trial (RAPPORT) Investigators. *Circulation*. 1998 Aug 25;98(8):734-41. [PubMed](#)

Bulkley BH, Roberts WC. Steroid therapy during acute myocardial infarction. A cause of delayed healing and of ventricular aneurysm. *Am J Med*. 1974 Feb;56(2):244-50. [PubMed](#)

Bulut S, Aengevaeren WR, Luijten HJ, Verheugt FW. Successful out-of-hospital cardiopulmonary resuscitation: what is the optimal in-hospital treatment strategy. *Resuscitation*. 2000 Oct;47(2):155-61. [PubMed](#)

Cannon CP, Braunwald E, McCabe CH, Rader DJ, Rouleau JL, Belder R, Joyal SV, Hill KA, Pfeffer MA, Skene AM, Pravastatin or Atorvastatin Evaluation and Infection Therapy-Thrombolysis in Myocardial Infarction 2. Intensive versus moderate lipid lowering with statins after acute coronary syndromes. *N Engl J Med*. 2004 Apr 8;350(15):1495-504. [PubMed](#)

Cannon CP, Steinberg BA, Murphy SA, Mega JL, Braunwald E. Meta-analysis of cardiovascular outcomes trials comparing intensive versus moderate statin therapy. *J Am Coll Cardiol*. 2006 Aug 1;48(3):438-45. [PubMed](#)

Cantor WJ, Fitchett D, Borgundvaag B, Ducas J, Heffernan M, Cohen EA, Morrison LJ, Langer A, Dzavik V, Mehta SR, Lazzam C, Schwartz B, Casanova A, Goodman SG, TRANSFER-AMI Trial Investigators. Routine early angioplasty after fibrinolysis for acute myocardial infarction. *N Engl J Med*. 2009 Jun 25;360(26):2705-18. [PubMed](#)

Caracciolo EA, Davis KB, Sopko G, Kaiser GC, Corley SD, Schaff H, Taylor HA, Chaitman BR. Comparison of surgical and medical group survival in patients with left main coronary artery disease. Long-term CASS experience. *Circulation*. 1995 May 1;91(9):2325-34. [PubMed](#)

Chareonthaitawee P, Gibbons RJ, Roberts RS, Christian TF, Burns R, Yusuf S. The impact of time to thrombolytic treatment on outcome in patients with acute myocardial infarction. For the CORE investigators (Collaborative Organisation for RheothRx Evaluation). *Heart*. 2000 Aug;84(2):142-8. [PubMed](#)

Chen EW, Canto JG, Parsons LS, Peterson ED, Littrell KA, Every NR, Gibson CM, Hochman JS, Ohman EM, Cheeks M, Barron HV, Investigators in the National Registry of Myocardial Infarction 2. Relation between hospital intra-aortic balloon counterpulsation volume and mortality in acute myocardial infarction complicated by cardiogenic shock. *Circulation*. 2003 Aug 26;108(8):951-7. [PubMed](#)

Chen ZM, Jiang LX, Chen YP, Xie JX, Pan HC, Peto R, Collins R, Liu LS, COMMIT (ClopIdogrel and Metoprolol in Myocardial Infarction Trial) collaborative . Addition of clopidogrel to aspirin in 45,852 patients with acute myocardial infarction: randomised placebo-controlled trial. *Lancet*. 2005 Nov 5;366(9497):1607-21. [PubMed](#)

Chen ZM, Pan HC, Chen YP, Peto R, Collins R, Jiang LX, Xie JX, Liu LS. Early intravenous then oral metoprolol in 45,852 patients with acute myocardial infarction: randomised placebo-controlled trial. *Lancet*. 2005 Nov 5;366(9497):1622-32. [PubMed](#)

Cholesterol Treatment Trialists' (CTT) Collaboration, Baigent C, Blackwell L, Emberson J, Holland LE, Reith C, Bhala N, Peto R, Barnes EH, Keech A, Simes J, Collins R. Efficacy and safety of more intensive lowering of LDL cholesterol: a meta-analysis of data from 170,000 participants in 26 randomised trials. *Lancet*. 2010 Nov 13;376(9753):1670-81. [PubMed](#)

Coleman EA, Parry C, Chalmers S, Min SJ. The care transitions intervention: results of a randomized controlled trial. *Arch Intern Med*. 2006 Sep 25;166(17):1822-8. [PubMed](#)

Collet JP, Montalescot G, Le May M, Borentain M, Gershlick A. Percutaneous coronary intervention after fibrinolysis: a multiple meta-analyses approach according to the type of strategy. *J Am Coll Cardiol*. 2006 Oct 3;48(7):1326-35. [PubMed](#)

Connolly SJ, Hallstrom AP, Cappato R, Schron EB, Kuck KH, Zipes DP, Greene HL, Boczor S, Domanski M, Follmann D, Gent M, Roberts RS. Meta-analysis of the implantable cardioverter defibrillator secondary prevention trials. AVID, CASH and CIDS studies. Antiarrhythmics vs Implantable Defibrillator study. Cardiac Arrest Study Hamburg. Canadian Implantable Defibrillator Study. *Eur Heart J*. 2000 Dec;21(24):2071-8. [PubMed](#)

CURRENT-OASIS 7 Investigators, Mehta SR, Bassand JP, Chrolavicius S, Diaz R, Eikelboom JW, Fox KA, Granger CB, Jolly S, Joyner CD, Rupprecht HJ, Widimsky P, Afzal R, Pogue J, Yusuf S. Dose comparisons of clopidogrel and aspirin in acute coronary syndromes. *N Engl J Med*. 2010 Sep 2;363(10):930-42. [PubMed](#)

Dalby M, Bouzamondo A, Lechat P, Montalescot G. Transfer for primary angioplasty versus immediate thrombolysis in acute myocardial infarction: a meta-analysis. *Circulation*. 2003 Oct 14;108(15):1809-14. [PubMed](#)

Dalrymple-Hay MJ, Langley SM, Sami SA, Haw M, Allen SM, Livesey SA, Lamb RK, Monro JL. Should coronary artery bypass grafting be performed at the same time as repair of a post-infarct ventricular septal defect. *Eur J Cardiothorac Surg*. 1998 Mar;13(3):286-92. [PubMed](#)

Dawood N, Vaccarino V, Reid KJ, Spertus JA, Hamid N, Parashar S, PREMIER Registry Investigators. Predictors of smoking cessation after a myocardial infarction: the role of institutional smoking cessation programs in improving success. *Arch Intern Med*. 2008 Oct 13;168(18):1961-7. [PubMed](#)

De Luca G, Bellandi F, Huber K, Noc M, Petronio AS, Amtz HR, Maioli M, Gabriel HM, Zorman S, DE Carlo M, Rakowski T, Gyongyosi M, Dudek D. Early glycoprotein IIb-IIIa inhibitors in primary angioplasty-abciximab long-term results (EGYPT-ALT) cooperation: individual patient's data meta-analysis. *J Thromb Haemost*. 2011 Dec;9(12):2361-70. [PubMed](#)

de Winter RJ, Verouden NJ, Wellens HJ, Wilde AA, Interventional Cardiology Group of the Academic Medical Center. A new ECG sign of proximal LAD occlusion. *N Engl J Med*. 2008 Nov 6;359(19):2071-3. [PubMed](#)

Di Mario C, Dudek D, Piscione F, Mielecki W, Savonitto S, Murena E, Dimopoulos K, Manari A, Gaspardone A, Ochala A, Zmudka K, Bolognese L, Steg PG, Flather M, CARESS-in-AMI (Combined Abciximab RE-teplase Stent Study in Acute Myocardial) Investigators. Immediate angioplasty versus standard therapy with rescue angioplasty after thrombolysis in the Combined Abciximab REteplase Stent Study in Acute Myocardial Infarction (CARESS-in-AMI): an open, prospective, randomised, multicentre trial. *Lancet*. 2008 Feb 16;371(9612):559-68. [PubMed](#)

Dieker HJ, Liem SS, El Aidi H, van Grunsven P, Aengevaeren WR, Brouwer MA, Verheugt FW. Pre-hospital triage for primary angioplasty.



direct referral to the intervention center versus interhospital transport. *JACC Cardiovasc Interv.* 2010 Jul;3(7):705-11. [PubMed](#)

Diercks DB, Kontos MC, Chen AY, Pollack CV Jr, Wiviott SD, Rumsfeld JS, Magid DJ, Gibler WB, Cannon CP, Peterson ED, Roe MT. Utilization and impact of pre-hospital electrocardiograms for patients with acute ST-segment elevation myocardial infarction: data from the NCDR (National Cardiovascular Data Registry) ACTION [trunc]. *J Am Coll Cardiol.* 2009 Jan 13;53(2):161-6. [PubMed](#)

D'Souza SP, Mamas MA, Fraser DG, Fath-Ordoubadi F. Routine early coronary angioplasty versus ischaemia-guided angioplasty after thrombolysis in acute ST-elevation myocardial infarction: a meta-analysis. *Eur Heart J.* 2011 Apr;32(8):972-82. [PubMed](#)

Dyke CM, Bhatia D, Lorenz TJ, Marso SP, Tardiff BE, Hogeboom C, Harrington RA. Immediate coronary artery bypass surgery after platelet inhibition with eptifibatide: results from PURSUIT. Platelet Glycoprotein IIb/IIIa in Unstable Angina: Receptor Suppression Using Integrilin Therapy. *Ann Thorac Surg.* 2000 Sep;70(3):866-71; discussion 871-2. [PubMed](#)

Early effects of tissue-type plasminogen activator added to conventional therapy on the culprit coronary lesion in patients presenting with ischemic cardiac pain at rest. Results of the Thrombolysis in Myocardial Ischemia (TIMI IIIA) Trial. *Circulation.* 1993 Jan;87(1):38-52. [PubMed](#)

Ebrahimi R, Dyke C, Mehran R, Manoukian SV, Feit F, Cox DA, Gersh BJ, Ohman EM, White HD, Moses JW, Ware JH, Lincoff AM, Stone GW. Outcomes following pre-operative clopidogrel administration in patients with acute coronary syndromes undergoing coronary artery bypass surgery: the ACUTY (Acute Catheterization and Urgent Intervention Triage strategY) trial. *J Am Coll Cardiol.* 2009 May 26;53(21):1965-72. [PubMed](#)

Effect of intravenous APSAC on mortality after acute myocardial infarction: preliminary report of a placebo-controlled clinical trial. AIMS Trial Study Group. *Lancet.* 1988 Mar 12;1(8585):545-9. [PubMed](#)

El Khoury C, Dubien PY, Mercier C, Belle L, Debaty G, Capel O, Perret T, Savary D, Serre P, Bonnefoy E, AGIR-2 Investigators. Prehospital high-dose tirofiban in patients undergoing primary percutaneous intervention. The AGIR-2 study. *Arch Cardiovasc Dis.* 2010 May;103(5):285-92. [PubMed](#)

Ellis SG, Armstrong P, Betriu A, Brodie B, Hermann H, Montalescot G, Neumann FJ, Smith JJ, Topol E, Facilitated Intervention with Enhanced Reperfusion Speed to Stop Events. Facilitated percutaneous coronary intervention versus primary percutaneous coronary intervention: design and rationale of the Facilitated Intervention with Enhanced Reperfusion Speed to Stop Events (FINESSE) trial. *Am Heart J.* 2004 Apr;147(4):E16. [PubMed](#)

Ellis SG, Tendera M, de Belder MA, van Boven AJ, Widimsky P, Andersen HR, Betriu A, Savonitto S, Adamus J, Peruga JZ, Hamankiewicz M, Pluta W, Oldroyd K, Ecollan P, Janssens L, Armstrong P, Brodie BR, Hermann HC, Montalescot G, Neumann FJ, Effron MB, Barnathan ES, Topol EJ, FINESSE Investigators. 1-year survival in a randomized trial of facilitated reperfusion: results from the FINESSE (Facilitated Intervention with Enhanced Reperfusion Speed to Stop Events) trial. *JACC Cardiovasc Interv.* 2009 Oct;2(10):909-16. [PubMed](#)

Epstein AE, DiMarco JP, Ellenbogen KA, Estes NA 3rd, Freedman RA, Gettes LS, Gillinov AM, Gregoratos G, Hammill SC, Hayes DL, Hlatky MA, Newby LK, Page RL, Schoenfeld MH, Silka MJ, Stevenson LW, Sweeney MO, Smith SC Jr, Jacobs AK, Adams CD, Anderson JL, Buller CE, Creager MA, Ettinger SM, Faxon DP, Halperin JL, Hiratzka LF, Hunt SA, Krumholz HM, Kushner FG, Lytle BW, Nishimura RA, Ornato JP, Page RL, Riegel B, Tarkington LG, Yancy CW, American College of Cardiology/American Heart Association Task Force on Practice, American Association for Thoracic Surgery, Society of Thoracic Surgeons. ACC/AHA/HRS 2008 guidelines for device-based therapy of cardiac rhythm abnormalities: a report of the American College of Cardiology/American Heart Association task force [trunc]. *J Am Coll Cardiol.* 2008 May 27;51(21):e1-62. [527 references] [PubMed](#)

Erne P, Schoenenberger AW, Burckhardt D, Zuber M, Kiowski W, Buser PT, Dubach P, Resink TJ, Pfisterer M. Effects of percutaneous coronary interventions in silent ischemia after myocardial infarction: the SWISSI II randomized controlled trial. *JAMA.* 2007 May 9;297(18):1985-91. [PubMed](#)

Faxon DP, Eikelboom JW, Berger PB, Holmes DR, Bhatt DL, Moliterno DJ, Becker RC, Angiolillo DJ. Consensus document: antithrombotic therapy in patients with atrial fibrillation undergoing coronary stenting. A North-American perspective. *Thromb Haemost.* 2011

Fernandez-Aviles F, Alonso JJ, Castro-Beiras A, Vazquez N, Blanco J, Alonso-Briales J, Lopez-Mesa J, Fernandez-Vazquez F, Calvo I, Martinez-Elbal L, San Roman JA, Ramos B. Routine invasive strategy within 24 hours of thrombolysis versus ischaemia-guided conservative approach for acute myocardial infarction with ST-segment elevation (GRACIA-1): a randomised controlled trial. *Lancet*. 2004 Sep 18-24;364(9439):1045-53. [PubMed](#)

Fibrinolytic Therapy Trialists' (FFT) Collaborative Group. Indications for fibrinolytic therapy in suspected acute myocardial infarction: collaborative overview of early mortality and major morbidity results from all randomised trials of more than 1000 patients. *Lancet*. 1994 Feb 5;343(8893):311-22. [58 references] [PubMed](#)

Freemantle N, Cleland J, Young P, Mason J, Harrison J. Beta blockade after myocardial infarction: systematic review and meta regression analysis. *BMJ*. 1999 Jun 26;318(7200):1730-7. [PubMed](#)

French JK, Feldman HA, Assmann SF, Sanborn T, Palmeri ST, Miller D, Boland J, Buller CE, Steingart R, Sleeper LA, Hochman JS. Influence of thrombolytic therapy, with or without intra-aortic balloon counterpulsation, on 12-month survival in the SHOCK trial. *Am Heart J*. 2003 Nov;146(5):804-10. [PubMed](#)

Garot P, Lefevre T, Eltchaninoff H, Morice MC, Tamion F, Abry B, Lesault PF, Le Tarnec JY, Pouges C, Margenet A, Monchi M, Laurent I, Dumas P, Garot J, Louvard Y. Six-month outcome of emergency percutaneous coronary intervention in resuscitated patients after cardiac arrest complicating ST-elevation myocardial infarction. *Circulation*. 2007 Mar 20;115(11):1354-62. [PubMed](#)

Gershlick AH, Stephens-Lloyd A, Hughes S, Abrams KR, Stevens SE, Uren NG, de Belder A, Davis J, Pitt M, Banning A, Baumbach A, Shiu MF, Schofield P, Dawkins KD, Henderson RA, Oldroyd KG, Wilcox R, REACT Trial Investigators. Rescue angioplasty after failed thrombolytic therapy for acute myocardial infarction. *N Engl J Med*. 2005 Dec 29;353(26):2758-68. [PubMed](#)

Gibson CM, Cannon CP, Murphy SA, Marble SJ, Barron HV, Braunwald E, TIMI Study Group. Relationship of the TIMI myocardial perfusion grades, flow grades, frame count, and percutaneous coronary intervention to long-term outcomes after thrombolytic administration in acute myocardial infarction. *Circulation*. 2002 Apr 23;105(16):1909-13. [PubMed](#)

Gibson CM, Karha J, Murphy SA, James D, Morrow DA, Cannon CP, Giugliano RP, Antman EM, Braunwald E, TIMI Study Group. Early and long-term clinical outcomes associated with reinfarction following fibrinolytic administration in the Thrombolysis in Myocardial Infarction trials. *J Am Coll Cardiol*. 2003 Jul 2;42(1):7-16. [PubMed](#)

Gibson CM, Murphy SA, Montalescot G, Morrow DA, Ardissino D, Cohen M, Gulba DC, Kracoff OH, Lewis BS, Roguin N, Antman EM, Braunwald E, ExTRACT-TIMI 25 Investigators. Percutaneous coronary intervention in patients receiving enoxaparin or unfractionated heparin after fibrinolytic therapy for ST-segment elevation myocardial infarction in the ExTRACT-TIMI 25 trial. *J Am Coll Cardiol*. 2007 Jun 12;49(23):2238-46. [PubMed](#)

Gibson CM, Murphy SA, Rizzo MJ, Ryan KA, Marble SJ, McCabe CH, Cannon CP, Van de Werf F, Braunwald E. Relationship between TIMI frame count and clinical outcomes after thrombolytic administration. Thrombolysis In Myocardial Infarction (TIMI) Study Group. *Circulation*. 1999 Apr 20;99(15):1945-50. [PubMed](#)

Gierlotka M, Gasior M, Wilczek K, Hawranek M, Szkodzinski J, Paczek P, Lekston A, Kalarus Z, Zembala M, Polonski L. Reperfusion by primary percutaneous coronary intervention in patients with ST-segment elevation myocardial infarction within 12 to 24 hours of the onset of symptoms (from a prospective national observational study [PL-ACS]). *Am J Cardiol*. 2011 Feb 15;107(4):501-8. [PubMed](#)

GISSI-3: effects of lisinopril and transdermal glyceryl trinitrate singly and together on 6-week mortality and ventricular function after acute myocardial infarction. Gruppo Italiano per lo Studio della Sopravvivenza nell'infarto Miocardico. *Lancet*. 1994 May 7;343(8906):1115-22. [PubMed](#)

Goel K, Lennon RJ, Tilbury RT, Squires RW, Thomas RJ. Impact of cardiac rehabilitation on mortality and cardiovascular events after percutaneous coronary intervention in the community. *Circulation*. 2011 May 31;123(21):2344-52. [PubMed](#)

Gorjup V, Radsel P, Kocjancic ST, Erzen D, Noc M. Acute ST-elevation myocardial infarction after successful cardiopulmonary resuscitation. *Resuscitation*. 2007 Mar;72(3):379-85. [PubMed](#)

Grines CL, Bonow RO, Casey DE Jr, Gardner TJ, Lockhart PB, Moliterno DJ, O'Gara P, Whitlow P, American Heart Association, American College of Cardiology, Society for Cardiovascular Angiography and Interventions, American College of Surgeons, American Dental Association, American College of Physicians. Prevention of premature discontinuation of dual antiplatelet therapy in patients with coronary artery stents: a science advisory from the American Heart Association, American College of Cardiology, Society for Cardiovascular Angiography [trunc]. *Circulation*. 2007 Feb 13;115(6):813-8. [36 references] [PubMed](#)

Grzybowski M, Clements EA, Parsons L, Welch R, Tintinalli AT, Ross MA, Zalenski RJ. Mortality benefit of immediate revascularization of acute ST-segment elevation myocardial infarction in patients with contraindications to thrombolytic therapy: a propensity analysis. *JAMA*. 2003 Oct 8;290(14):1891-8. [PubMed](#)

Gupta M, Chang WC, Van de Werf F, Granger CB, Midodji W, Barbash G, Pehrson K, Oto A, Toutouzas P, Jansky P, Armstrong PW. International differences in in-hospital revascularization and outcomes following acute myocardial infarction: a multilevel analysis of patients in ASSENT-2. *Eur Heart J*. 2003 Sep;24(18):1640-50. [PubMed](#)

Hannan EL, Samadashvili Z, Walford G, Holmes DR Jr, Jacobs AK, Stamato NJ, Venditti FJ, Sharma S, King SB 3rd. Culprit vessel percutaneous coronary intervention versus multivessel and staged percutaneous coronary intervention for ST-segment elevation myocardial infarction patients with multivessel disease. *JACC Cardiovasc Interv*. 2010 Jan;3(1):22-31. [PubMed](#)

Held C, Asenblad N, Bassand JP, Becker RC, Cannon CP, Claeys MJ, Harrington RA, Horrow J, Husted S, James SK, Mahaffey KW, Nicolau JC, Scirica BM, Storey RF, Vintila M, Ycas J, Wallentin L. Ticagrelor versus clopidogrel in patients with acute coronary syndromes undergoing coronary artery bypass surgery: results from the PLATO (Platelet Inhibition and Patient Outcomes) trial. *J Am Coll Cardiol*. 2011 Feb 8;57(6):672-84. [PubMed](#)

Henry TD, Sharkey SW, Burke MN, Chavez IJ, Graham KJ, Henry CR, Lips DL, Madison JD, Menssen KM, Mooney MR, Newell MC, Pedersen WR, Poulouse AK, Traverse JH, Unger BT, Wang YL, Larson DM. A regional system to provide timely access to percutaneous coronary intervention for ST-elevation myocardial infarction. *Circulation*. 2007 Aug 14;116(7):721-8. [PubMed](#)

Hochman JS, Buller CE, Sleeper LA, Boland J, Dzavik V, Sanborn TA, Godfrey E, White HD, Lim J, LeJemtel T. Cardiogenic shock complicating acute myocardial infarction--etiologies, management and outcome: a report from the SHOCK Trial Registry. Should we emergently revascularize Occluded Coronaries for cardiogenic shock. *J Am Coll Cardiol*. 2000 Sep;36(3 Suppl A):1063-70. [PubMed](#)

Hochman JS, Lamas GA, Buller CE, Dzavik V, Reynolds HR, Abramsky SJ, Forman S, Ruzyllo W, Maggioni AP, White H, Sadowski Z, Carvalho AC, Rankin JM, Renkin JP, Steg PG, Mascette AM, Sopko G, Pfisterer ME, Leor J, Fridrich V, Mark DB, Knatterud GL, Occluded Artery Trial Investigators. Coronary intervention for persistent occlusion after myocardial infarction. *N Engl J Med*. 2006 Dec 7;355(23):2395-407. [PubMed](#)

Hochman JS, Sleeper LA, Webb JG, Dzavik V, Buller CE, Aylward P, Col J, White HD, SHOCK Investigators. Early revascularization and long-term survival in cardiogenic shock complicating acute myocardial infarction. *JAMA*. 2006 Jun 7;295(21):2511-5. [PubMed](#)

Hochman JS, Sleeper LA, Webb JG, Sanborn TA, White HD, Talley JD, Buller CE, Jacobs AK, Slater JN, Col J, McKinlay SM, LeJemtel TH. Early revascularization in acute myocardial infarction complicated by cardiogenic shock. SHOCK Investigators. Should We Emergently Revascularize Occluded Coronaries for Cardiogenic Shock. *N Engl J Med*. 1999 Aug 26;341(9):625-34. [PubMed](#)

Hochman JS, Sleeper LA, White HD, Dzavik V, Wong SC, Menon V, Webb JG, Steingart R, Picard MH, Menegus MA, Boland J, Sanborn T, Buller CE, Modur S, Forman R, Desvigne-Nickens P, Jacobs AK, Slater JN, LeJemtel TH, SHOCK Investigators. One-year survival



following early revascularization for cardiogenic shock. JAMA. 2001 Jan 10;285(2):190-2. [PubMed](#)

Hohnloser SH, Kuck KH, Dorian P, Roberts RS, Hampton JR, Hatala R, Fain E, Gent M, Connolly SJ, DINAMIT Investigators. Prophylactic use of an implantable cardioverter-defibrillator after acute myocardial infarction. N Engl J Med. 2004 Dec 9;351(24):2481-8. [PubMed](#)

Hosmane VR, Mustafa NG, Reddy VK, Reese CL 4th, DiSabatino A, Kolm P, Hopkins JT, Weintraub WS, Rahman E. Survival and neurologic recovery in patients with ST-segment elevation myocardial infarction resuscitated from cardiac arrest. J Am Coll Cardiol. 2009 Feb 3;53(5):409-15. [PubMed](#)

Hypothermia after Cardiac Arrest Study Group. Mild therapeutic hypothermia to improve the neurologic outcome after cardiac arrest. N Engl J Med 2002 Feb 21;346(8):549-56. [PubMed](#)

Indications for ACE inhibitors in the early treatment of acute myocardial infarction: systematic overview of individual data from 100,000 patients in randomized trials. ACE Inhibitor Myocardial Infarction Collaborative Group. Circulation. 1998 Jun 9;97(22):2202-12. [PubMed](#)

Ioannidis JP, Katritsis DG. Percutaneous coronary intervention for late reperfusion after myocardial infarction in stable patients. Am Heart J. 2007 Dec;154(6):1065-71. [PubMed](#)

ISIS-4: a randomised factorial trial assessing early oral captopril, oral mononitrate, and intravenous magnesium sulphate in 58,050 patients with suspected acute myocardial infarction. ISIS-4 (Fourth International Study of Infarct Survival) Collabora. Lancet. 1995 Mar 18;345(8951):669-85. [PubMed](#)

Iversen A, Galatius S, Jensen JS. The optimal route of administration of the glycoprotein IIb/IIIa receptor antagonist abciximab during percutaneous coronary intervention; intravenous versus intracoronary. Curr Cardiol Rev. 2008 Nov;4(4):293-9. [PubMed](#)

Jack BW, Chetty VK, Anthony D, Greenwald JL, Sanchez GM, Johnson AE, Forsythe SR, O'Donnell JK, Paasche-Orlow MK, Manasseh C, Martin S, Culpepper L. A reengineered hospital discharge program to decrease rehospitalization: a randomized trial. Ann Intern Med. 2009 Feb 3;150(3):178-87. [PubMed](#)

Jacob M, Smedira N, Blackstone E, Williams S, Cho L. Effect of timing of chronic preoperative aspirin discontinuation on morbidity and mortality in coronary artery bypass surgery. Circulation. 2011 Feb 15;123(6):577-83. [PubMed](#)

Jeremias A, Sylvia B, Bridges J, Kirtane AJ, Bigelow B, Pinto DS, Ho KK, Cohen DJ, Garcia LA, Cutlip DE, Carrozza JP Jr. Stent thrombosis after successful sirolimus-eluting stent implantation. Circulation. 2004 Apr 27;109(16):1930-2. [PubMed](#)

Jollis JG, Roettig ML, Aluko AO, Anstrom KJ, Applegate RJ, Babb JD, Berger PB, Bohle DJ, Fletcher SM, Garvey JL, Hathaway WR, Hoekstra JW, Kelly RV, Maddox WT Jr, Shiber JR, Valeri FS, Watling BA, Wilson BH, Granger CB, Reperfusion of Acute Myocardial Infarction in North Carolina Emergency. Implementation of a statewide system for coronary reperfusion for ST-segment elevation myocardial infarction. JAMA. 2007 Nov 28;298(20):2371-80. [PubMed](#)

Jolly SS, Pogue J, Haladyn K, Peters RJ, Fox KA, Avezum A, Gersh BJ, Rupprecht HJ, Yusuf S, Mehta SR. Effects of aspirin dose on ischaemic events and bleeding after percutaneous coronary intervention: insights from the PCI-CURE study. Eur Heart J. 2009 Apr;30(8):900-7. [PubMed](#)

Jong GP, Ma T, Chou P, Shyu MY, Tseng WK, Chang TC. Reciprocal changes in 12-lead electrocardiography can predict left main coronary artery lesion in patients with acute myocardial infarction. Int Heart J. 2006 Jan;47(1):13-20. [PubMed](#)

Kahn JK, Glazier S, Swor R, Savas V, O'Neill WW. Primary coronary angioplasty for acute myocardial infarction complicated by out-of-hospital cardiac arrest. Am J Cardiol. 1995 May 15;75(15):1069-70. [PubMed](#)

Kakkar AK, Moustapha A, Hanley HG, Weiss M, Caldito G, Misra P, Reddy PC, Tandon N. Comparison of intracoronary vs. intravenous administration of abciximab in coronary stenting. *Catheter Cardiovasc Interv.* 2004 Jan;61(1):31-4. [PubMed](#)

Kaluza GL, Joseph J, Lee JR, Raizner ME, Raizner AE. Catastrophic outcomes of noncardiac surgery soon after coronary stenting. *J Am Coll Cardiol.* 2000 Apr;35(5):1288-94. [PubMed](#)

Keelan PC, Bunch TJ, White RD, Packer DL, Holmes DR Jr. Early direct coronary angioplasty in survivors of out-of-hospital cardiac arrest. *Am J Cardiol.* 2003 Jun 15;91(12):1461-3, A6. [PubMed](#)

Keeley EC, Boura JA, Grines CL. Comparison of primary and facilitated percutaneous coronary interventions for ST-elevation myocardial infarction: quantitative review of randomised trials. *Lancet.* 2006 Feb 18;367(9510):579-88. [28 references] [PubMed](#)

Keeley EC, Boura JA, Grines CL. Primary angioplasty versus intravenous thrombolytic therapy for acute myocardial infarction: a quantitative review of 23 randomised trials. *Lancet.* 2003 Jan 4;361(9351):13-20. [41 references] [PubMed](#)

Kern KB, Rahman O. Emergent percutaneous coronary intervention for resuscitated victims of out-of-hospital cardiac arrest. *Catheter Cardiovasc Interv.* 2010 Mar 1;75(4):616-24. [26 references] [PubMed](#)

Kim JH, Newby LK, Clare RM, Shaw LK, Lodge AJ, Smith PK, Jolicœur EM, Rao SV, Becker RC, Mark DB, Granger CB. Clopidogrel use and bleeding after coronary artery bypass graft surgery. *Am Heart J.* 2008 Nov;156(5):886-92. [PubMed](#)

Kober L, Torp-Pedersen C, Carlsen JE, Bagger H, Eliassen P, Lyngborg K, Videbaek J, Cole DS, Auclert L, Pauly NC. A clinical trial of the angiotensin-converting-enzyme inhibitor trandolapril in patients with left ventricular dysfunction after myocardial infarction. Trandolapril Cardiac Evaluation (TRACE) Study Group. *N Engl J Med.* 1995 Dec 21;333(25):1670-6. [PubMed](#)

Lappe JM, Muhlestein JB, Lappe DL, Badger RS, Bair TL, Brockman R, French TK, Hofmann LC, Home BD, Kralick-Goldberg S, Nicponski N, Orton JA, Pearson RR, Renlund DG, Rimmasch H, Roberts C, Anderson JL. Improvements in 1-year cardiovascular clinical outcomes associated with a hospital-based discharge medication program. *Ann Intern Med.* 2004 Sep 21;141(6):446-53. [PubMed](#)

Late Assessment of Thrombolytic Efficacy (LATE) study with alteplase 6-24 hours after onset of acute myocardial infarction. *Lancet.* 1993 Sep 25;342(8874):759-66. [PubMed](#)

Le May MR, So DY, Dionne R, Glover CA, Froeschl MP, Wells GA, Davies RF, Sherrard HL, Maloney J, Marquis JF, O'Brien ER, Trickett J, Poirier P, Ryan SC, Ha A, Joseph PG, Labinaz M. A citywide protocol for primary PCI in ST-segment elevation myocardial infarction. *N Engl J Med.* 2008 Jan 17;358(3):231-40. [PubMed](#)

Leon AS, Franklin BA, Costa F, Balady GJ, Berra KA, Stewart KJ, Thompson PD, Williams MA, Lauer MS. Cardiac rehabilitation and secondary prevention of coronary heart disease: an AHA scientific statement from the Council on Clinical Cardiology and the Council on Nutrition, Physical Activity, and Metabolism, in collaboration with AACVPR. *Circulation.* 2005 Jan 25;111(3):369-76. [PubMed](#)

Leppo JA, O'Brien J, Rothendler JA, Getchell JD, Lee VW. Dipyridamole-thallium-201 scintigraphy in the prediction of future cardiac events after acute myocardial infarction. *N Engl J Med.* 1984 Apr 19;310(16):1014-8. [PubMed](#)

Lip GY, Huber K, Andreotti F, Arnesen H, Airaksinen JK, Cuisset T, Kirchhof P, Marin F, Consensus Document of European Society of Cardiology Working Group on Thrombosis. Antithrombotic management of atrial fibrillation patients presenting with acute coronary syndrome and/or undergoing coronary stenting: executive summary [trunc]. *Eur Heart J.* 2010 Jun;31(11):1311-8. [PubMed](#)

Madsen JK, Grande P, Saunamäki K, Thayssen P, Kassiss E, Eriksen U, Rasmussen K, Haunso S, Nielsen TT, Haghfelt T, Fritz-Hansen P, Hjelm E, Paulsen PK, Alstrup P, Arendrup H, Niebuhr-Jørgensen U, Andersen LI. Danish multicenter randomized study of invasive versus

conservative treatment in patients with inducible ischemia after thrombolysis in acute myocardial infarction (DANAMI). DANish trial in Acute Myocardial Infarction. *Circulation*. 1997 Aug 5;96(3):748-55. [PubMed](#)

Maggioni AP, Fabbri G. VALIANT (VALsartan In Acute myocardial iNfarcTion) trial. *Expert Opin Pharmacother*. 2005 Mar;6(3):507-12. [PubMed](#)

Maioli M, Bellandi F, Leoncini M, Toso A, Dabizzi RP. Randomized early versus late abciximab in acute myocardial infarction treated with primary coronary intervention (RELAX-AMI Trial). *J Am Coll Cardiol*. 2007 Apr 10;49(14):1517-24. [PubMed](#)

Maltais S, Perrault LP, Do QB. Effect of clopidogrel on bleeding and transfusions after off-pump coronary artery bypass graft surgery: impact of discontinuation prior to surgery. *Eur J Cardiothorac Surg*. 2008 Jul;34(1):127-31. [PubMed](#)

Marcusohn E, Roguin A, Sebbag A, Aronson D, Dragu R, Amikam S, Boulus M, Grenadier E, Kerner A, Nikolsky E, Markiewicz W, Hammerman H, Kapeliovich M. Primary percutaneous coronary intervention after out-of-hospital cardiac arrest: patients and outcomes. *Isr Med Assoc J*. 2007 Apr;9(4):257-9. [PubMed](#)

McNamara RL, Herrin J, Wang Y, Curtis JP, Bradley EH, Magid DJ, Rathore SS, Nallamothu BK, Peterson ED, Blaney ME, Frederick P, Krumholz HM. Impact of delay in door-to-needle time on mortality in patients with ST-segment elevation myocardial infarction. *Am J Cardiol*. 2007 Oct 15;100(8):1227-32. [PubMed](#)

Mehilli J, Kastrati A, Schulz S, Frungel S, Nekolla SG, Moshage W, Dotzer F, Huber K, Pache J, Dirschinger J, Seyfarth M, Martinoff S, Schwaiger M, Schomig A, Bavarian Reperfusion Alternatives Evaluation-3 (BRAVE-3) Study Investigators. Abciximab in patients with acute ST-segment-elevation myocardial infarction undergoing primary percutaneous coronary intervention after clopidogrel loading: a randomized double-blind trial. *Circulation*. 2009 Apr 14;119(14):1933-40. [PubMed](#)

Mehta SR, Tanguay JF, Eikelboom JW, Jolly SS, Joyner CD, Granger CB, Faxon DP, Rupprecht HJ, Budaj A, Avezum A, Widimsky P, Steg PG, Bassand JP, Montalescot G, Macaya C, Di Pasquale G, Niemela K, Ajani AE, White HD, Chrolavicius S, Gao P, Fox KA, Yusuf S, CURRENT-OASIS 7 trial investigators. Double-dose versus standard-dose clopidogrel and high-dose versus low-dose aspirin in individuals undergoing percutaneous coronary intervention for acute coronary syndromes (CURRENT-OASIS 7): a randomised factorial trial. *Lancet*. 2010 Oct 9;376(9748):1233-43. [PubMed](#)

Menon V, Webb JG, Hillis LD, Sleeper LA, Abboud R, Dzavik V, Slater JN, Forman R, Monrad ES, Talley JD, Hochman JS. Outcome and profile of ventricular septal rupture with cardiogenic shock after myocardial infarction: a report from the SHOCK Trial Registry. Should we emergently revascularize Occluded Coronaries in cardiogenic shock. *J Am Coll Cardiol*. 2000 Sep;36(3 Suppl A):1110-6. [PubMed](#)

Milavetz JJ, Giebel DW, Christian TF, Schwartz RS, Holmes DR Jr, Gibbons RJ. Time to therapy and salvage in myocardial infarction. *J Am Coll Cardiol*. 1998 May;31(6):1246-51. [PubMed](#)

Montalescot G, Barragan P, Wittenberg O, Ecollan P, Elhadad S, Villain P, Boulenc JM, Morice MC, Maillard L, Pansieri M, Choussat R, Pinton P. Platelet glycoprotein IIb/IIIa inhibition with coronary stenting for acute myocardial infarction. *N Engl J Med*. 2001 Jun 21;344(25):1895-903. [PubMed](#)

Montalescot G, Borentain M, Payot L, Collet JP, Thomas D. Early vs late administration of glycoprotein IIb/IIIa inhibitors in primary percutaneous coronary intervention of acute ST-segment elevation myocardial infarction: a meta-analysis. *JAMA*. 2004 Jul 21;292(3):362-6. [PubMed](#)

Montalescot G, Wiviott SD, Braunwald E, Murphy SA, Gibson CM, McCabe CH, Antman EM, TRITON-TIMI 38 investigators. Prasugrel compared with clopidogrel in patients undergoing percutaneous coronary intervention for ST-elevation myocardial infarction (TRITON-TIMI 38): double-blind, randomised controlled trial. *Lancet*. 2009 Feb 28;373(9665):723-31. [PubMed](#)

Morrow DA, Antman EM, Charlesworth A, Cairns R, Murphy SA, de Lemos JA, Giugliano RP, McCabe CH, Braunwald E. TIMI risk score

for ST-elevation myocardial infarction: A convenient, bedside, clinical score for risk assessment at presentation: An intravenous nPA for treatment of infarcting myocardium early II trial substudy. *Circulation*. 2000 Oct 24;102(17):2031-7. [PubMed](#)

Moss AJ, Zareba W, Hall WJ, Klein H, Wilber DJ, Cannom DS, Daubert JP, Higgins SL, Brown MW, Andrews ML. Prophylactic implantation of a defibrillator in patients with myocardial infarction and reduced ejection fraction. *N Engl J Med*. 2002 Mar 21;346(12):877-83. [PubMed](#)

Nallamothu BK, Bates ER. Percutaneous coronary intervention versus fibrinolytic therapy in acute myocardial infarction: is timing (almost) everything. *Am J Cardiol*. 2003 Oct 1;92(7):824-6. [PubMed](#)

Nasser M, Kapeliovich M, Markiewicz W. Late thrombosis of sirolimus-eluting stents following noncardiac surgery. *Catheter Cardiovasc Interv*. 2005 Aug;65(4):516-9. [PubMed](#)

Naylor M, Brooten D, Jones R, Lavizzo-Mourey R, Mezey M, Pauly M. Comprehensive discharge planning for the hospitalized elderly. A randomized clinical trial. *Ann Intern Med*. 1994 Jun 15;120(12):999-1006. [PubMed](#)

Newby LK, Rutsch WR, Califf RM, Simoons ML, Aylward PE, Armstrong PW, Woodlief LH, Lee KL, Topol EJ, Van de Werf F. Time from symptom onset to treatment and outcomes after thrombolytic therapy. GUSTO-1 Investigators. *J Am Coll Cardiol*. 1996 Jun;27(7):1646-55. [PubMed](#)

Nichol G, Aufderheide TP, Eigel B, Neumar RW, Lurie KG, Bufalino VJ, Callaway CW, Menon V, Bass RR, Abella BS, Sayre M, Dougherty CM, Racht EM, Kleinman ME, O'Connor RE, Reilly JP, Ossmann EW, Peterson E, American Heart Association Emergency Cardiovascular Care Committee, Council on Arteriosclerosis, Thrombosis, and Vascular Biology, Council on Cardiopulmonary, Critical Care, Perioperative and Resuscitation, Council on Cardiovascular Nursing, Council on Clinical Cardiology, Advocacy Committee, Council on Quality of Care and Outcomes Research. Regional systems of care for out-of-hospital cardiac arrest: A policy statement from the American Heart Association. *Circulation*. 2010 Feb 9;121(5):709-29. [PubMed](#)

Nielsen PH, Terkelsen CJ, Nielsen TT, Thuesen L, Krusell LR, Thayssen P, Kelbaek H, Abildgaard U, Villadsen AB, Andersen HR, Maeng M, Danami-2 Investigators. System delay and timing of intervention in acute myocardial infarction (from the Danish Acute Myocardial Infarction-2 [DANAMI-2] trial). *Am J Cardiol*. 2011 Sep 15;108(6):776-81. [PubMed](#)

Nijjer SS, Watson G, Athanasiou T, Malik IS. Safety of clopidogrel being continued until the time of coronary artery bypass grafting in patients with acute coronary syndrome: a meta-analysis of 34 studies. *Eur Heart J*. 2011 Dec;32(23):2970-88. [PubMed](#)

Nordmann AJ, Hengstler P, Harr T, Young J, Bucher HC. Clinical outcomes of primary stenting versus balloon angioplasty in patients with myocardial infarction: a meta-analysis of randomized controlled trials. *Am J Med*. 2004 Feb 15;116(4):253-62. [31 references] [PubMed](#)

Ohman EM, Nanas J, Stomel RJ, Leeser MA, Nielsen DW, O'Dea D, Rogers FJ, Harber D, Hudson MP, Fraulo E, Shaw LK, Lee KL, TACTICS Trial. Thrombolysis and counterpulsation to improve survival in myocardial infarction complicated by hypotension and suspected cardiogenic shock or heart failure: results of the TACTICS Trial. *J Thromb Thrombolysis*. 2005 Feb;19(1):33-9. [PubMed](#)

Park DW, Park SW, Park KH, Lee BK, Kim YH, Lee CW, Hong MK, Kim JJ, Park SJ. Frequency of and risk factors for stent thrombosis after drug-eluting stent implantation during long-term follow-up. *Am J Cardiol*. 2006 Aug 1;98(3):352-6. [PubMed](#)

Patti G, Barcsi G, Orlic D, Mangiacapra F, Colonna G, Pasceri V, Barbato E, Merkely B, Edes I, Ostojic M, Wijns W, Di Sciascio G. Outcome comparison of 600- and 300-mg loading doses of clopidogrel in patients undergoing primary percutaneous coronary intervention for ST-segment elevation myocardial infarction [trunc] . *J Am Coll Cardiol*. 2011 Oct 4;58(15):1592-9. [PubMed](#)

Peberdy MA, Callaway CW, Neumar RW, Geocadin RG, Zimmerman JL, Donnino M, Gabrielli A, Silvers SM, Zaritsky AL, Merchant R, Vanden Hoek TL, Kronick SL. Part 9: post-cardiac arrest care: 2010 American Heart Association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation*. 2010 Nov 2;122(18 Suppl 3):S768-86. [PubMed](#)

Pfeiffer MA, Braunwald E, Moye LA, Basta L, Brown EJ Jr, Cuddy TE, Davis BR, Geltman EM, Goldman S, Flaker GC, et al. Effect of captopril on mortality and morbidity in patients with left ventricular dysfunction after myocardial infarction. Results of the survival and ventricular enlargement trial. The SAVE Investigators. *N Engl J Med*. 1992 Sep 3;327(10):669-77. [PubMed](#)

Pfeiffer MA, Greaves SC, Arnold JM, Glynn RJ, LaMotte FS, Lee RT, Menapace FJ Jr, Rapaport E, Ridker PM, Rouleau JL, Solomon SD, Hennekens CH. Early versus delayed angiotensin-converting enzyme inhibition therapy in acute myocardial infarction. The healing and early afterload reducing therapy trial. *Circulation*. 1997 Jun 17;95(12):2643-51. [PubMed](#)

Pfeiffer MA, McMurray JJ, Velazquez EJ, Rouleau JL, Kober L, Maggioni AP, Solomon SD, Swedberg K, Van de Werf F, White H, Leimberger JD, Henis M, Edwards S, Zelenkofske S, Sellers MA, Califf RM. Valsartan, captopril, or both in myocardial infarction complicated by heart failure, left ventricular dysfunction, or both. *N Engl J Med*. 2003 Nov 13;349(20):1893-906. [PubMed](#)

Pfisterer M, Brunner-La Rocca HP, Buser PT, Rickenbacher P, Hunziker P, Mueller C, Jeger R, Bader F, Osswald S, Kaiser C, BASKET-LATE Investigators. Late clinical events after clopidogrel discontinuation may limit the benefit of drug-eluting stents: an observational study of drug-eluting versus bare-metal stents. *J Am Coll Cardiol*. 2006 Dec 19;48(12):2584-91. [PubMed](#)

Pinto DS, Kirtane AJ, Nallamothu BK, Murphy SA, Cohen DJ, Laham RJ, Cutlip DE, Bates ER, Frederick PD, Miller DP, Carrozza JP Jr, Antman EM, Cannon CP, Gibson CM. Hospital delays in reperfusion for ST-elevation myocardial infarction: implications when selecting a reperfusion strategy. *Circulation*. 2006 Nov 7;114(19):2019-25. [PubMed](#)

Pitt B, Remme W, Zannad F, Neaton J, Martinez F, Roniker B, Bittman R, Hurley S, Kleinman J, Gatlin M. Eplerenone, a selective aldosterone blocker, in patients with left ventricular dysfunction after myocardial infarction. *N Engl J Med*. 2003 Apr 3;348(14):1309-21. [PubMed](#)

Pleskot M, Babu A, Hazukova R, Stritecky J, Bis J, Matejka J, Cermakova E. Out-of-hospital cardiac arrests in patients with acute ST elevation myocardial infarctions in the East Bohemian region over the period 2002-2004. *Cardiology*. 2008;109(1):41-51. [PubMed](#)

Quintero-Moran B, Moreno R, Villarreal S, Perez-Vizcayno MJ, Hernandez R, Conde C, Vazquez P, Alfonso F, Banuelos C, Escaned J, Fernandez-Ortiz A, Azcona L, Macaya C. Percutaneous coronary intervention for cardiac arrest secondary to ST-elevation acute myocardial infarction. Influence of immediate paramedical/medical assistance on clinical outcome. *J Invasive Cardiol*. 2006 Jun;18(6):269-72. [PubMed](#)

Randomised trial of intravenous atenolol among 16,027 cases of suspected acute myocardial infarction: ISIS-1. ISIS-1 (First International Study of Infarct Survival Collaborative Group). *Lancet*. 1986 Jul 12;2(8498):57-66. [PubMed](#)

Randomised trial of intravenous streptokinase, oral aspirin, both, or neither among 17,187 cases of suspected acute myocardial infarction: ISIS-2. ISIS-2 (Second International Study of Infarct Survival) Collaborative Group. *Lancet*. 1988 Aug 13;2(8607):349-60. [PubMed](#)

Randomised trial of late thrombolysis in patients with suspected acute myocardial infarction. EMERAS (Estudio Multicentrico Estreptoquinasa Republicas de America del Sur) Collaborative Group. *Lancet*. 1993 Sep 25;342(8874):767-72. [PubMed](#)

Richling N, Herkner H, Holzer M, Riedmueller E, Sterz F, Schreiber W. Thrombolytic therapy vs primary percutaneous intervention after ventricular fibrillation cardiac arrest due to acute ST-segment elevation myocardial infarction and its effect on outcome. *Am J Emerg Med*. 2007 Jun;25(5):545-50. [PubMed](#)

Roberts R, Rogers WJ, Mueller HS, Lambrew CT, Diver DJ, Smith HC, Willerson JT, Knatterud GL, Forman S, Passamani E, et al. Immediate versus deferred beta-blockade following thrombolytic therapy in patients with acute myocardial infarction. Results of the Thrombolysis in Myocardial Infarction (TIMI) II-B Study. *Circulation*. 1991 Feb;83(2):422-37. [PubMed](#)

Rokos IC, French WJ, Koenig WJ, Stratton SJ, Nighswonger B, Strunk B, Jewell J, Mahmud E, Dunford JV, Hokanson J, Smith SW, Baran KW, Swor R, Berman A, Wilson BH, Aluko AO, Gross BW, Rostkyus PS, Salvucci A, Dev V, McNally B, Manoukian SV, King SB 3rd.



Integration of pre-hospital electrocardiograms and ST-elevation myocardial infarction receiving center (SRC) networks: impact on Door-to-Balloon times across 10 independent regions. *JACC Cardiovasc Interv.* 2009 Apr;2(4):339-46. [PubMed](#)

Romagnoli E, Burzotta F, Trani C, Mazzari MA, Biondi-Zoccai GG, De Vita M, Giannico F, Niccoli G, Prati F, Rebuzzi AG, Mongiardo R, Crea F. Angiographic evaluation of the effect of intracoronary abciximab administration in patients undergoing urgent PCI. *Int J Cardiol.* 2005 Dec 7;105(3):250-5. [PubMed](#)

Ross AM, Molhoek P, Lundergan C, Knudtson M, Draoui Y, Regalado L, Le Louer V, Bigonzi F, Schwartz W, de Jong E, Coyne K, HART II Investigators. Randomized comparison of enoxaparin, a low-molecular-weight heparin, with unfractionated heparin adjunctive to recombinant tissue plasminogen activator thrombolysis and aspirin: second trial of Heparin and Aspirin Reperfusion Therapy (HART II). *Circulation.* 2001 Aug 7;104(6):648-52. [PubMed](#)

Rossi P, Bolognese L. Comparison of intravenous urokinase plus heparin versus heparin alone in acute myocardial infarction. Urochinas per via Sistemica nell'Infarto Miocardico (USIM) Collaborative Group. *Am J Cardiol.* 1991 Sep 1;68(6):585-92. [PubMed](#)

Sabatine MS, Cannon CP, Gibson CM, Lopez-Sendon JL, Montalescot G, Theroux P, Claeys MJ, Cools F, Hill KA, Skene AM, McCabe CH, Braunwald E. Addition of clopidogrel to aspirin and fibrinolytic therapy for myocardial infarction with ST-segment elevation. *N Engl J Med.* 2005 Mar 24;352(12):1179-89. [PubMed](#)

Sanborn TA, Sleeper LA, Bates ER, Jacobs AK, Boland J, French JK, Dens J, Dzavik V, Palmeri ST, Webb JG, Goldberger M, Hochman JS. Impact of thrombolysis, intra-aortic balloon pump counterpulsation, and their combination in cardiogenic shock complicating acute myocardial infarction: a report from the SHOCK Trial Registry. SHow we emergently revascularize [trunc]. *J Am Coll Cardiol.* 2000 Sep;36(3 Suppl A):1123-9. [PubMed](#)

Sardella G, Mancone M, Bucciarelli-Ducci C, Agati L, Scardala R, Carbone I, Francone M, Di Roma A, Benedetti G, Conti G, Fedele F. Thrombus aspiration during primary percutaneous coronary intervention improves myocardial reperfusion and reduces infarct size: the EXPIRA (thrombectomy with export catheter in infarct-related artery during primary percutaneous coronary [trunc]. *J Am Coll Cardiol.* 2009 Jan 27;53(4):309-15. [PubMed](#)

Schomig A, Mehilli J, Antoniucci D, Ndrepepa G, Markwardt C, Di Pede F, Nekolla SG, Schlotterbeck K, Schuhlen H, Pache J, Seyfarth M, Martinoff S, Benzer W, Schmitt C, Dirschinger J, Schwaiger M, Kastrati A, Beyond 12 hours Reperfusion Alternative Evaluation (BRAVE-2) Trial Investigators. Mechanical reperfusion in patients with acute myocardial infarction presenting more than 12 hours from symptom onset: a randomized controlled trial. *JAMA.* 2005 Jun 15;293(23):2865-72. [PubMed](#)

Schomig A, Neumann FJ, Kastrati A, Schuhlen H, Blasini R, Hadamitzky M, Walter H, Zitzmann-Roth EM, Richardt G, Alt E, Schnitt C, Ulm K. A randomized comparison of antiplatelet and anticoagulant therapy after the placement of coronary-artery stents. *N Engl J Med.* 1996 Apr 25;334(17):1084-9. [PubMed](#)

Serebruany VL, Steinhubl SR, Berger PB, Malinin AI, Baggish JS, Bhatt DL, Topol EJ. Analysis of risk of bleeding complications after different doses of aspirin in 192,036 patients enrolled in 31 randomized controlled trials. *Am J Cardiol.* 2005 May 15;95(10):1218-22. [PubMed](#)

Shah AM, Pfeffer MA, Hartley LH, Moye LA, Gersh BJ, Rutherford JD, Lamas GA, Rouleau JL, Braunwald E, Solomon SD. Risk of all-cause mortality, recurrent myocardial infarction, and heart failure hospitalization associated with smoking status following myocardial infarction with left ventricular dysfunction. *Am J Cardiol.* 2010 Oct 1;106(7):911-6. [PubMed](#)

Shim JK, Choi YS, Oh YJ, Bang SO, Yoo KJ, Kwak YL. Effects of preoperative aspirin and clopidogrel therapy on perioperative blood loss and blood transfusion requirements in patients undergoing off-pump coronary artery bypass graft surgery. *J Thorac Cardiovasc Surg.* 2007 Jul;134(1):59-64. [PubMed](#)

Siebel J, Kuck KH. Implantable cardioverter defibrillator compared with antiarrhythmic drug treatment in cardiac arrest survivors (the Cardiac Arrest Study Hamburg). *Am Heart J.* 1994 Apr;127(4 Pt 2):1139-44. [PubMed](#)

Silverman HS, Pfeifer MP. Relation between use of anti-inflammatory agents and left ventricular free wall rupture during acute myocardial infarction. *Am J Cardiol.* 1987 Feb 1;59(4):363-4. [PubMed](#)

Sjaauw KD, Engstrom AE, Vis MM, van der Schaaf RJ, Baan J Jr, Koch KT, de Winter RJ, Piek JJ, Tijssen JG, Henriques JP. A systematic review and meta-analysis of intra-aortic balloon pump therapy in ST-elevation myocardial infarction: should we change the guidelines. *Eur Heart J.* 2009 Feb;30(4):459-68. [43 references] [PubMed](#)

Slater J, Brown RJ, Antonelli TA, Menon V, Boland J, Col J, Dzavik V, Greenberg M, Menegus M, Connery C, Hochman JS. Cardiogenic shock due to cardiac free-wall rupture or tamponade after acute myocardial infarction: a report from the SHOCK Trial Registry. Should we emergently revascularize occluded coronaries for cardiogenic shock. *J Am Coll Cardiol.* 2000 Sep;36(3 Suppl A):1117-22. [PubMed](#)

Smith SC Jr, Benjamin EJ, Bonow RO, Braun LT, Creager MA, Franklin BA, Gibbons RJ, Grundy SM, Hiratzka LF, Jones DW, Lloyd-Jones DM, Minissian M, Mosca L, Peterson ED, Sacco RL, Spertus J, Stein JH, Taubert KA. AHA/ACC secondary prevention and risk reduction therapy for patients with coronary and other atherosclerotic vascular disease: 2011 update: a guideline from the American Heart Association and American College of Cardiology Foundation [trunc]. *J Am Coll Cardiol.* 2011 Nov 29;58(23):2432-46. [PubMed](#)

Sorensen JT, Terkelsen CJ, Norgaard BL, Trautner S, Hansen TM, Botker HE, Lassen JF, Andersen HR. Urban and rural implementation of pre-hospital diagnosis and direct referral for primary percutaneous coronary intervention in patients with acute ST-elevation myocardial infarction. *Eur Heart J.* 2011 Feb;32(4):430-6. [PubMed](#)

Spaulding CM, Joly LM, Rosenberg A, Monchi M, Weber SN, Dhainaut JF, Carli P. Immediate coronary angiography in survivors of out-of-hospital cardiac arrest. *N Engl J Med.* 1997 Jun 5;336(23):1629-33. [PubMed](#)

Spertus JA, Kettelkamp R, Vance C, Decker C, Jones PG, Rumsfeld JS, Messenger JC, Khanal S, Peterson ED, Bach RG, Krumholz HM, Cohen DJ. Prevalence, predictors, and outcomes of premature discontinuation of thienopyridine therapy after drug-eluting stent placement: results from the PREMIER registry. *Circulation.* 2006 Jun 20;113(24):2803-9. [PubMed](#)

Steg PG, Dabbous OH, Feldman LJ, Cohen-Solal A, Aumont MC, Lopez-Sendon J, Budaj A, Goldberg RJ, Klein W, Anderson FA Jr, Global Registry of Acute Coronary Events Investigators. Determinants and prognostic impact of heart failure complicating acute coronary syndromes: observations from the Global Registry of Acute Coronary Events (GRACE). *Circulation.* 2004 Feb 3;109(4):494-9. [PubMed](#)

Steg PG, James S, Harrington RA, Ardissino D, Becker RC, Cannon CP, Emanuelsson H, Finkelstein A, Husted S, Katus H, Kilhamn J, Olofsson S, Storey RF, Weaver WD, Wallentin L, PLATO Study Group. Ticagrelor versus clopidogrel in patients with ST-elevation acute coronary syndromes intended for reperfusion with primary percutaneous coronary intervention: A Platelet Inhibition and Patient Outcomes (PLATO) trial subgroup analysis. *Circulation.* 2010 Nov 23;122(21):2131-41. [PubMed](#)

Steg PG, Kerner A, Van de Werf F, Lopez-Sendon J, Gore JM, Fitzgerald G, Feldman LJ, Anderson FA, Avezum A, Global Registry of Acute Coronary Events (GRACE) Investigators. Impact of in-hospital revascularization on survival in patients with non-ST-elevation acute coronary syndrome and congestive heart failure. *Circulation.* 2008 Sep 9;118(11):1163-71. [PubMed](#)

Steg PG, Thuaire C, Himbert D, Carrie D, Champagne S, Coisne D, Khalife K, Cazaux P, Logeart D, Slama M, Spaulding C, Cohen A, Tirouvanziam A, Montely JM, Rodriguez RM, Garbarz E, Wijns W, Durand-Zaleski I, Porcher R, Brucker L, Chevret S, Chastang C, DECOPI Investigators. DECOPI (DEsobstruction CORonaire en Post-Infarctus): a randomized multi-centre trial of occluded artery angioplasty after acute myocardial infarction. *Eur Heart J.* 2004 Dec;25(24):2187-94. [PubMed](#)

Steinbeck G, Andresen D, Seidl K, Brachmann J, Hoffmann E, Wojciechowski D, Kornacewicz-Jach Z, Sredniawa B, Lupkovics G, Hofgartner F, Lubinski A, Rosenqvist M, Habets A, Wegscheider K, Senges J, IRIS Investigators. Defibrillator implantation early after myocardial infarction. *N Engl J Med.* 2009 Oct 8;361(15):1427-36. [PubMed](#)

Steinhubl SR, Bhatt DL, Brennan DM, Montalescot G, Hankey GJ, Eikelboom JW, Berger PB, Topol EJ, CHARISMA Investigators. Aspirin

to prevent cardiovascular disease: the association of aspirin dose and clopidogrel with thrombosis and bleeding. *Ann Intern Med.* 2009 Mar 17;150(6):379-86. [PubMed](#)

Stone GW, Grines CL, Cox DA, Garcia E, Tcheng JE, Griffin JJ, Guagliumi G, Stuckey T, Turco M, Carroll JD, Rutherford BD, Lansky AJ. Comparison of angioplasty with stenting, with or without abciximab, in acute myocardial infarction. *N Engl J Med.* 2002 Mar 28;346(13):957-66. [PubMed](#)

Stone GW, Maehara A, Witzenbichler B, Godlewski J, Parise H, Dambrink JH, Ochala A, Carlton TW, Cristea E, Wolff SD, Brener SJ, Chowdhary S, El-Omar M, Neunteufl T, Metzger DC, Karwoski T, Dizon JM, Mehran R, Gibson CM, INFUSE-AMI Investigators. Intracoronary abciximab and aspiration thrombectomy in patients with large anterior myocardial infarction: the INFUSE-AMI randomized trial. *JAMA.* 2012 May 2;307(17):1817-26. [PubMed](#)

Stone GW, Witzenbichler B, Guagliumi G, Peruga JZ, Brodie BR, Dudek D, Kornowski R, Hartmann F, Gersh BJ, Pocock SJ, Dangas G, Wong SC, Kirtane AJ, Parise H, Mehran R. Bivalirudin during primary PCI in acute myocardial infarction. *N Engl J Med.* 2008 May 22;358(21):2218-30. [30 references] [PubMed](#)

Suaya JA, Stason WB, Ades PA, Normand SL, Shepard DS. Cardiac rehabilitation and survival in older coronary patients. *J Am Coll Cardiol.* 2009 Jun 30;54(1):25-33. [PubMed](#)

Sutton AG, Campbell PG, Graham R, Price DJ, Gray JC, Grech ED, Hall JA, Harcombe AA, Wright RA, Smith RH, Murphy JJ, Shyam-Sundar A, Stewart MJ, Davies A, Linker NJ, de Belder MA. A randomized trial of rescue angioplasty versus a conservative approach for failed fibrinolysis in ST-segment elevation myocardial infarction: the Middlesbrough Early Revascularization to Limit Infarction (MERLIN) trial. *J Am Coll Cardiol.* 2004 Jul 21;44(2):287-96. [PubMed](#)

Sutton AG, Campbell PG, Price DJ, Grech ED, Hall JA, Davies A, Stewart MJ, de Belder MA. Failure of thrombolysis by streptokinase: detection with a simple electrocardiographic method. *Heart.* 2000 Aug;84(2):149-56. [PubMed](#)

Tavakoli R, Weber A, Vogt P, Brunner HP, Pretre R, Turina M. Surgical management of acute mitral valve regurgitation due to post-infarction papillary muscle rupture. *J Heart Valve Dis.* 2002 Jan;11(1):20-5; discussion 26. [PubMed](#)

Taylor RS, Brown A, Ebrahim S, Jolliffe J, Noorani H, Rees K, Skidmore B, Stone JA, Thompson DR, Oldridge N. Exercise-based rehabilitation for patients with coronary heart disease: systematic review and meta-analysis of randomized controlled trials. *Am J Med.* 2004 May 15;116(10):682-92. [75 references] [PubMed](#)

ten Berg JM, van't Hof AW, Dill T, Heestermans T, van Werkum JW, Mosterd A, van Houwelingen G, Koopmans PC, Stella PR, Boersma E, Hamm C, On-TIME 2 Study Group. Effect of early, pre-hospital initiation of high bolus dose tirofiban in patients with ST-segment elevation myocardial infarction on short- and long-term clinical outcome. *J Am Coll Cardiol.* 2010 Jun 1;55(22):2446-55. [PubMed](#)

Theroux P, Waters DD, Halphen C, Debaisieux JC, Mizgala HF. Prognostic value of exercise testing soon after myocardial infarction. *N Engl J Med.* 1979 Aug 16;301(7):341-5. [PubMed](#)

Thompson CR, Buller CE, Sleeper LA, Antonelli TA, Webb JG, Jaber WA, Abel JG, Hochman JS. Cardiogenic shock due to acute severe mitral regurgitation complicating acute myocardial infarction: a report from the SHOCK Trial Registry. Should we use emergently revascularize Occluded Coronaries in cardiogenic shock. *J Am Coll Cardiol.* 2000 Sep;36(3 Suppl A):1104-9. [PubMed](#)

Thomson CC, Rigotti NA. Hospital- and clinic-based smoking cessation interventions for smokers with cardiovascular disease. *Prog Cardiovasc Dis.* 2003 May-Jun;45(6):459-79. [174 references] [PubMed](#)

Thune JJ, Hoefsten DE, Lindholm MG, Mortensen LS, Andersen HR, Nielsen TT, Kober L, Kelbaek H, Danish Multicenter Randomized Study on Fibrinolytic Therapy Versus Acute Coronary. Simple risk stratification at admission to identify patients with reduced mortality from primary angioplasty. *Circulation.* 2005 Sep 27;112(13):2017-21. [PubMed](#)



Toma M, Buller CE, Westerhout CM, Fu Y, O'Neill WW, Holmes DR Jr, Hamm CW, Granger CB, Armstrong PW, APEX-AMI Investigators. Non-culprit coronary artery percutaneous coronary intervention during acute ST-segment elevation myocardial infarction: insights from the APEX-AMI trial. *Eur Heart J*. 2010 Jul;31(14):1701-7. [PubMed](#)

Valgimigli M, Campo G, Percoco G, Bolognese L, Vassanelli C, Colangelo S, de Cesare N, Rodriguez AE, Ferrario M, Moreno R, Piva T, Sheiban I, Pasquetto G, Prati F, Nazzaro MS, Parrinello G, Ferrari R, Multicentre Evaluation of Single High-Dose Bolus Tirofiban vs Abciximab With Sirolimus-Eluting Stent. Comparison of angioplasty with infusion of tirofiban or abciximab and with implantation of sirolimus-eluting or uncoated stents for acute myocardial infarction: the MULTISTRATEGY randomized trial. *JAMA*. 2008 Apr 16;299(15):1788-99. [PubMed](#)

Vandvik PO, Lincoff AM, Gore JM, Gutterman DD, Sonnenberg FA, Alonso-Coello P, Akl EA, Lansberg MG, Guyatt GH, Spencer FA. Primary and secondary prevention of cardiovascular disease: antithrombotic therapy and prevention of thrombosis, 9th ed: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest*. 2012 Feb;141(2 Suppl):e637S-68S. [101 references] [PubMed](#)

Van't Hof AW, Ten Berg J, Heestermaars T, Dill T, Funck RC, van Werkum W, Dambrink JH, Suryapranata H, van Houwelingen G, Ottervanger JP, Stella P, Giannitsis E, Hamm C, Ongoing Tirofiban In Myocardial infarction Evaluation (On-TIME) 2 study group. Prehospital initiation of tirofiban in patients with ST-elevation myocardial infarction undergoing primary angioplasty (On-TIME 2): a multicentre, double-blind, randomised controlled trial. *Lancet*. 2008 Aug 16;372(9638):537-46. [PubMed](#)

Villella A, Maggioni AP, Villella M, Giordano A, Turazza FM, Santoro E, Franzosi MG. Prognostic significance of maximal exercise testing after myocardial infarction treated with thrombolytic agents: the GISSI-2 data-base. Gruppo Italiano per lo Studio della Sopravvivenza Nell'Infarto. *Lancet*. 1995 Aug 26;346(8974):523-9. [PubMed](#)

Vlaar PJ, Mahmoud KD, Holmes DR Jr, van Valkenhoef G, Hillege HL, van der Horst IC, Zijlstra F, de Smet BJ. Culprit vessel only versus multivessel and staged percutaneous coronary intervention for multivessel disease in patients presenting with ST-segment elevation myocardial infarction: a pairwise and network meta-analysis. *J Am Coll Cardiol*. 2011 Aug 9;58(7):692-703. [PubMed](#)

Vlaar PJ, Svilaas T, van der Horst IC, Diercks GF, Fokkema ML, de Smet BJ, van den Heuvel AF, Anthonio RL, Jessurun GA, Tan ES, Suurmeijer AJ, Zijlstra F. Cardiac death and reinfarction after 1 year in the Thrombus Aspiration during Percutaneous coronary intervention in Acute myocardial infarction Study (TAPAS): a 1-year follow-up study. *Lancet*. 2008 Jun 7;371(9628):1915-20. [PubMed](#)

Werling M, Thoren AB, Axelsson C, Herlitz J. Treatment and outcome in post-resuscitation care after out-of-hospital cardiac arrest when a modern therapeutic approach was introduced. *Resuscitation*. 2007 Apr;73(1):40-5. [PubMed](#)

Wever EF, Hauer RN, van Capelle FL, Tijssen JG, Crijns HJ, Algra A, Wiesfeld AC, Bakker PF, Robles de Medina EO. Randomized study of implantable defibrillator as first-choice therapy versus conventional strategy in postinfarct sudden death survivors. *Circulation*. 1995 Apr 15;91(8):2195-203. [PubMed](#)

White HD. Systems of care: need for hub-and-spoke systems for both primary and systematic percutaneous coronary intervention after fibrinolysis. *Circulation*. 2008 Jul 15;118(3):219-22. [23 references] [PubMed](#)

Wijeyesundera HC, Vijayaraghavan R, Nallamotheu BK, Foody JM, Krumholz HM, Phillips CO, Kashani A, You JJ, Tu JV, Ko DT. Rescue angioplasty or repeat fibrinolysis after failed fibrinolytic therapy for ST-segment myocardial infarction: a meta-analysis of randomized trials. *J Am Coll Cardiol*. 2007 Jan 30;49(4):422-30. [PubMed](#)

Wilson K, Gibson N, Willan A, Cook D. Effect of smoking cessation on mortality after myocardial infarction: meta-analysis of cohort studies. *Arch Intern Med*. 2000 Apr 10;160(7):939-44. [PubMed](#)

Wilson SH, Bell MR, Rihal CS, Bailey KR, Holmes DR Jr, Berger PB. Infarct artery reocclusion after primary angioplasty, stent placement, and thrombolytic therapy for acute myocardial infarction. *Am Heart J*. 2001 May;141(5):704-10. [PubMed](#)

Wiviott SD, Braunwald E, McCabe CH, Montalescot G, Ruzyllo W, Gottlieb S, Neumann FJ, Ardissino D, De Servi S, Murphy SA, Riesmeyer J, Weerakkody G, Gibson CM, Antman EM, TRITON-TIMI 38 Investigators. Prasugrel versus clopidogrel in patients with acute coronary syndromes. *N Engl J Med*. 2007 Nov 15;357(20):2001-15. [PubMed](#)

Wohrle J, Grebe OC, Nusser T, Al-Khayer E, Schaible S, Kochs M, Hombach V, Hoher M. Reduction of major adverse cardiac events with intracoronary compared with intravenous bolus application of abciximab in patients with acute myocardial infarction or unstable angina undergoing coronary angioplasty. *Circulation*. 2003 Apr 15;107(14):1840-3. [PubMed](#)

Woo YJ, Grand T, Valettas N. Off-pump coronary artery bypass grafting attenuates postoperative bleeding associated with preoperative clopidogrel administration. *Heart Surg Forum*. 2003;6(5):282-5. [PubMed](#)

Wu AH, Parsons L, Every NR, Bates ER, Second National Registry of Myocardial Infarction. Hospital outcomes in patients presenting with congestive heart failure complicating acute myocardial infarction: a report from the Second National Registry of Myocardial Infarction (NRMII-2). *J Am Coll Cardiol*. 2002 Oct 16;40(8):1389-94. [PubMed](#)

You JJ, Singer DE, Howard PA, Lane DA, Eckman MH, Fang MC, Hylek EM, Schulman S, Go AS, Hughes M, Spencer FA, Manning WJ, Halperin JL, Lip GY. Antithrombotic therapy for atrial fibrillation: antithrombotic therapy and prevention of thrombosis, 9th ed: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest*. 2012 Feb;141(2 Suppl):e531S-75S. [175 references] [PubMed](#)

Young W, Rewa G, Goodman SG, Jaglal SB, Cash L, Lefkowitz C, Coyte PC. Evaluation of a community-based inner-city disease management program for postmyocardial infarction patients: a randomized controlled trial. *CMAJ*. 2003 Oct 28;169(9):905-10. [PubMed](#)

Yusuf S, Mehta SR, Chrolavicius S, Afzal R, Pogue J, Granger CB, Budaj A, Peters RJ, Bassand JP, Wallentin L, Joyner C, Fox KA, OASIS-6 Trial Group. Effects of fondaparinux on mortality and reinfarction in patients with acute ST-segment elevation myocardial infarction: the OASIS-6 randomized trial. *JAMA*. 2006 Apr 5;295(13):1519-30. [PubMed](#)

Zahn R, Schuster S, Schiele R, Seidl K, Voigtlander T, Meyer J, Hauptmann KE, Gottwik M, Berg G, Kunz T, Gieseler U, Jakob M, Senges J. Comparison of primary angioplasty with conservative therapy in patients with acute myocardial infarction and contraindications for thrombolytic therapy. Maximal Individual Therapy in Acute Myocardial Infarction (MITRA) Study Group. *Catheter Cardiovasc Interv*. 1999 Feb;46(2):127-33. [PubMed](#)

Zhu MM, Feit A, Chadow H, Alam M, Kwan T, Clark LT. Primary stent implantation compared with primary balloon angioplasty for acute myocardial infarction: a meta-analysis of randomized clinical trials. *Am J Cardiol*. 2001 Aug 1;88(3):297-301. [PubMed](#)

Zijlstra F, Hoorntje JC, de Boer MJ, Reiffers S, Miedema K, Ottervanger JP, van 't Hof AW, Suryapranata H. Long-term benefit of primary angioplasty as compared with thrombolytic therapy for acute myocardial infarction. *N Engl J Med*. 1999 Nov 4;341(19):1413-9. [PubMed](#)

## Type of Evidence Supporting the Recommendations

The type of supporting evidence is identified and graded for each recommendation (see the "Major Recommendations" field).

## Benefits/Harms of Implementing the Guideline Recommendations

### Potential Benefits

- Decreased morbidity and mortality due to ST-elevation myocardial infarction (STEMI)
- Effective secondary prevention of myocardial infarction and other cardiovascular events

## Potential Harms

- Potential complications of primary percutaneous coronary intervention (PCI) include problems with the arterial access site; adverse reactions to volume loading, contrast medium, and antithrombotic medications; technical complications; and reperfusion events.
- Nonfatal and fatal bleeding can occur in patients treated with fibrinolytic therapy, aspirin, clopidogrel, and/or vitamin K antagonists.
- Although "false positives" are a concern when emergency medical services (EMS) personnel and/or emergency physicians are allowed to activate the cardiac catheterization laboratory, the rate of false activations is relatively low (approximately 15%) and is more than balanced by earlier treatment times for the majority of patients for whom notification is appropriate.
- Until further data become available, it would seem prudent to weigh the possible increased risk of intracranial bleeding when the addition of ticagrelor to aspirin is considered in patients with prior stroke or transient ischemic attack.
- Glycoprotein IIb/IIIa inhibitors should be used with great caution, if at all, after full-dose fibrinolytic therapy, because this combination is associated with high rates of bleeding and intracranial hemorrhage (ICH), particularly in the elderly.
- Although pericarditis is not an absolute contraindication to anticoagulation, caution should be exercised because of the potential for hemorrhagic conversion.
- Caution is advised when using calcium channel blockers in patients with left ventricular systolic dysfunction.
- Heparin-induced thrombocytopenia (HIT), with and without thrombotic complications, can infrequently complicate the course of patients with acute coronary syndrome (ACS), particularly patients who previously have been exposed to heparin or who receive heparin over several hospital days. From 1% to 5% of all patients receiving heparin will develop HIT, and of these, 25% to 50% will develop thrombotic complications.

See Table 12 in the original guideline document for risks of selected medical therapies.

## Contraindications

### Contraindications

- Fondaparinux should not be given as the sole anticoagulant to patients referred for percutaneous coronary intervention (PCI) and is contraindicated for patients with a creatinine clearance less than 30 mL/min.
- Drug-eluting stents (DES) should not be used in primary PCI for patients with ST-elevation myocardial infarction (STEMI) who are unable to tolerate or comply with a prolonged course of dual antiplatelet therapy (DAPT) because of the increased risk of stent thrombosis with premature discontinuation of one or both agents.
- Avoid eptifibatide in patients on hemodialysis.
- Oral beta-blockers are contraindicated in patients with signs of heart failure (HF), evidence of a low output state, increased risk for cardiogenic shock.
- Refer to Table 12 in the original guideline document for contraindications/cautions of routine medical therapy used in STEMI.

### Contraindications and Cautions for Fibrinolytic Therapy in STEMI\*

#### *Absolute Contraindications*

- Any prior intracranial hemorrhage
- Known structural cerebral vascular lesion (e.g., arteriovenous malformation)
- Known malignant intracranial neoplasm (primary or metastatic)
- Ischemic stroke within 3 months (EXCEPT acute ischemic stroke within 4.5 h)
- Suspected aortic dissection
- Active bleeding or bleeding diathesis (excluding menses)
- Significant closed-head or facial trauma within 3 months
- Intracranial or intraspinal surgery within 2 months
- Severe uncontrolled hypertension (unresponsive to emergency therapy)
- For streptokinase, prior treatment within the previous 6 months

#### *Relative Contraindications*

- History of chronic, severe, poorly controlled hypertension
- Significant hypertension on presentation (systolic blood pressure >180 mm Hg or diastolic blood pressure >110 mm Hg)

- History of prior ischemic stroke >3 months
- Dementia
- Known intracranial pathology not covered in absolute contraindications
- Traumatic or prolonged (>10 min) cardiopulmonary resuscitation (CPR)
- Major surgery (<3 weeks)
- Recent (within 2 to 4 weeks) internal bleeding
- Noncompressible vascular punctures
- Pregnancy
- Active peptic ulcer
- Oral anticoagulant therapy

\*Viewed as advisory for clinical decision making and may not be all-inclusive or definitive.

## Qualifying Statements

### Qualifying Statements

- Because the American College of Cardiology Foundation (ACCF) and the American Heart Association (AHA) practice guidelines address patient populations (and healthcare providers) residing in North America, drugs that are not currently available in North America are discussed in the text without a specific class of recommendation (COR). For studies performed in large numbers of subjects outside North America, each writing committee reviews the potential influence of different practice patterns and patient populations on the treatment effect and relevance to the ACCF/AHA target population to determine whether the findings should inform a specific recommendation.
- The ACCF/AHA practice guidelines are intended to assist healthcare providers in clinical decision making by describing a range of generally acceptable approaches to the diagnosis, management, and prevention of specific diseases or conditions. The guidelines attempt to define practices that meet the needs of most patients in most circumstances. The ultimate judgment regarding care of a particular patient must be made by the healthcare provider and patient in light of all the circumstances presented by that patient. As a result, situations may arise for which deviations from these guidelines may be appropriate. Clinical decision making should involve consideration of the quality and availability of expertise in the area where care is provided. When these guidelines are used as the basis for regulatory or payer decisions, the goal should be improvement in quality of care. The Task Force recognizes that situations arise in which additional data are needed to inform patient care more effectively; these areas are identified within each respective guideline when appropriate.
- Prescribed courses of treatment in accordance with these recommendations are effective only if followed. Because lack of patient understanding and adherence may adversely affect outcomes, physicians and other healthcare providers should make every effort to engage the patient's active participation in prescribed medical regimens and lifestyles. In addition, patients should be informed of the risks, benefits, and alternatives to a particular treatment and should be involved in shared decision making whenever feasible, particularly for COR IIa and IIb, for which the benefit-to-risk ratio may be lower.

## Implementation of the Guideline

### Description of Implementation Strategy

An implementation strategy was not provided.

### Implementation Tools

Clinical Algorithm

Quick Reference Guides/Physician Guides

Slide Presentation

For information about availability, see the *Availability of Companion Documents* and *Patient Resources* fields below.

# Institute of Medicine (IOM) National Healthcare Quality Report Categories

## IOM Care Need

Getting Better

Living with Illness

## IOM Domain

Effectiveness

Patient-centeredness

Timeliness

## Identifying Information and Availability

### Bibliographic Source(s)

O'Gara PT, Kushner FG, Ascheim DD, Casey DE Jr, Chung MK, de Lemos JA, Ettinger SM, Fang JC, Fesmire FM, Franklin BA, Granger CB, Krumholz HM, Linderbaum JA, Morrow DA, Newby LK, Ornato JP, Ou N, Radford MJ, Tamis-Holland JE, Tommaso CL, Tracy CM, Woo YJ, Zhao DX. 2013 ACCF/AHA guideline for the management of ST-elevation myocardial infarction: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. J Am Coll Cardiol. 2013 Jan 29;61(4):e78-140. [656 references] [PubMed](#)

### Adaptation

Not applicable: The guideline was not adapted from another source.

### Date Released

1996 Nov 1 (revised 2013 Jan 29)

### Guideline Developer(s)

American College of Cardiology Foundation - Medical Specialty Society

American Heart Association - Professional Association

### Source(s) of Funding

The American College of Cardiology Foundation and the American Heart Association

### Guideline Committee

American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines

## Composition of Group That Authored the Guideline

*Writing Committee Members:* Patrick T. O'Gara, MD, FACC, FAHA (*Chair*); Frederick G. Kushner, MD, FACC, FAHA, FSCAI (*Vice Chair*); Deborah D. Ascheim, MD, FACC; Donald E. Casey, JR, MD, MPH, MBA, FACP, FAHA; Mina K. Chung, MD, FACC, FAHA; James A. de Lemos, MD, FACC; Steven M. Ettinger, MD, FACC; James C. Fang, MD, FACC, FAHA; Francis M. Fesmire, MD, FACEP; Barry A. Franklin, PhD, FAHA; Christopher B. Granger, MD, FACC, FAHA; Harlan M. Krumholz, MD, SM, FACC, FAHA; Jane A. Linderbaum, MS, CNP-BC; David A. Morrow, MD, MPH, FACC, FAHA; L. Kristin Newby, MD, MHS, FACC, FAHA; Joseph P. Ornato, MD, FACC, FAHA, FACP, FACEP; Narith Ou, PharmD; Martha J. Radford, MD, FACC, FAHA; Jacqueline E. Tamis-Holland, MD, FACC; Carl L. Tommaso, MD, FACC, FAHA, FSCAI; Cynthia M. Tracy, MD, FACC, FAHA; Y. Joseph Woo, MD, FACC, FAHA; David X. Zhao, MD, FACC

*ACCF/AHA Task Force Members:* Jeffrey L. Anderson, MD, FACC, FAHA (*Chair*); Alice K. Jacobs, MD, FACC, FAHA (*Immediate Past Chair*); Jonathan L. Halperin, MD, FACC, FAHA (*Chair-Elect*); Nancy M. Albert, PhD, CCNS, CCRN, FAHA; Ralph G. Brindis, MD, MPH, MACC; Mark A. Creager, MD, FACC, FAHA; David DeMets, PhD; Robert A. Guyton, MD, FACC, FAHA; Judith S. Hochman, MD, FACC, FAHA; Richard J. Kovacs, MD, FACC; Frederick G. Kushner, MD, FACC, FAHA\*; E. Magnus Ohman, MD, FACC; William G. Stevenson, MD, FACC, FAHA; Clyde W. Yancy, MD, FACC, FAHA\*

\*Former Task Force member during this writing effort.

## Financial Disclosures/Conflicts of Interest

The Task Force makes every effort to avoid actual, potential, or perceived conflicts of interest that may arise as a result of relationships with industry and other entities (RWI) among the members of the writing committee. All writing committee members and peer reviewers of the guideline are required to disclose all current healthcare related relationships, including those existing 12 months before initiation of the writing effort. In December 2009, the American College of Cardiology Foundation (ACCF) and American Heart Association (AHA) implemented a new RWI policy that requires the writing committee chair plus a minimum of 50% of the writing committee to have no *relevant* RWI (Appendix 1 in the original guideline document includes the ACCF/AHA definition of *relevance*). These statements are reviewed by the Task Force and all members during each conference call and/or meeting of the writing committee, and members provide updates as changes occur. All guideline recommendations require a confidential vote by the writing committee and must be approved by a consensus of the voting members. Members may not draft or vote on any text or recommendations pertaining to their RWI. Members who recused themselves from voting are indicated in the list of writing committee members, and specific section recusals are noted in Appendix 1 in the original guideline document. Authors' and peer reviewers' RWI pertinent to this guideline are disclosed in Appendixes 1 and 2, respectively in the original guideline document. In addition, to ensure complete transparency, writing committee members' comprehensive disclosure information—including RWI not pertinent to this document—is available as an online supplement. Comprehensive disclosure information for the Task Force is also available from the [American College of Cardiology Web site](#) . The work of the writing committees is supported exclusively by the ACCF and AHA without commercial support. Writing committee members volunteered their time for this activity.

## Guideline Endorser(s)

American College of Emergency Physicians - Medical Specialty Society

Society for Cardiovascular Angiography and Interventions - Medical Specialty Society

## Guideline Status

This is the current release of the guideline.

This guideline updates previous versions: Antman EM, Anbe DT, Armstrong PW, Bates ER, Green LA, Hand M, Hochman JS, Krumholz HM, Kushner FG, Lamas GA, Mullany CJ, Ornato JP, Pearle DL, Sloan MA, Smith SC Jr. ACC/AHA guidelines for the management of patients with ST-elevation myocardial infarction. A report of the Am Coll of Cardiol/American Heart Association Task Force on Practice Guidelines (Committee to revise the 1999 guidelines). Bethesda (MD): American College of Cardiology (ACC), American Heart Association (AHA); 2004. 211 p.

Canadian Cardiovascular Society, American Academy of Family Physicians, American College of Cardiology, American Heart Association, Antman EM, Hand M, Armstrong PW, Bates ER, Green LA, Halasyamani LK, Hochman JS, Krumholz HM, Lamas GA, Mullany CJ, Pearle



DL, Sloan MA, Smith SC Jr, Anbe DT, Kushner FG, Ornato JP, Pearle DL, Sloan MA, Jacobs AK, Adams CD, Anderson JL, Buller CE, Creager MA, Ettinger SM, Halperin JL, Hunt SA, Lytle BW, Nishimura R, Page RL, Riegel B, Tarkington LG, Yancy CW. 2007 focused update of the ACC/AHA 2004 guidelines for the management of patients with ST-elevation myocardial infarction: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. J Am Coll Cardiol 2008 Jan 15;51(2):210-47.

## Guideline Availability

Electronic copies: Available from the [Journal of the American College of Cardiology Web site](#)  and from the [Circulation Web site](#) .

Print copies: Available from the American College of Cardiology, 2400 N Street NW, Washington DC, 20037; (800) 253-4636 (US only).

## Availability of Companion Documents

The following are available:

- O'Gara PT, Kushner FG, Ascheim DD, Casey DE Jr, Chung MK, de Lemos JA, Ettinger SM, Fang JC, Fesmire FM, Franklin BA, Granger CB, Krumholz HM, Linderbaum JA, Morrow DA, Newby LK, Ornato JP, Ou N, Radford MJ, Tanis-Holland JE, Tommaso CL, Tracy CM, Woo YJ, Zhao DX. 2013 ACCF/AHA guideline for the management of ST-elevation myocardial infarction: executive summary. A report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. J Am Coll Cardiol 2013;61(4):485-510. Electronic copies: Available in Portable Document Format (PDF) from the [Journal of the American College of Cardiology \(JACC\) Web site](#) .
- 2013 ACCF/AHA guideline for the management of ST-elevation myocardial infarction. A report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. Slide set. 2013. 126 p. Electronic copies: Available from the [JACC Web site](#) .
- 2013 STEMI guideline data supplements. 2013. 17 p. Electronic copies: Available in PDF from the [JACC Web site](#) .
- Top ten things to know about the management of ST-elevation myocardial infarction. 2013. 1 p. Electronic copies: Available in PDF from the [American Heart Association \(AHA\) Web site](#) .
- Methodology manual and policies from the ACCF/AHA Task Force on Practice Guidelines. 2010 Jun. 88 p. Electronic copies: Available in Portable Document Format (PDF) from the [AHA Web site](#) .

## Patient Resources

None available

## NGC Status

The original NGC summary was completed by ECRI on June 30, 1998. The summary was updated by ECRI on September 2, 1999. This updated information was verified by the guideline developer on October 8, 1999. This summary was updated most recently on September 10, 2004. The updated information was verified by the guideline developer on February 23, 2005. This summary was updated on May 3, 2005 following the withdrawal of Bextra (valdecoxib) from the market and the release of heightened warnings for Celebrex (celecoxib) and other nonselective nonsteroidal anti-inflammatory drugs (NSAIDs). This summary was updated by ECRI on June 16, 2005, following the U.S. Food and Drug Administration advisory on COX-2 selective and non-selective non-steroidal anti-inflammatory drugs (NSAIDs). This summary was updated by ECRI on March 6, 2007 following the FDA advisory on Coumadin (warfarin sodium). This summary was updated by ECRI Institute on June 22, 2007 following the U.S. Food and Drug Administration (FDA) advisory on heparin sodium injection. This summary was updated by ECRI Institute on September 7, 2007 following the revised U.S. Food and Drug Administration (FDA) advisory on Coumadin (warfarin). This summary was updated by ECRI Institute on November 6, 2007, following the U.S. Food and Drug Administration advisory on Antidepressant drugs. This NGC summary was updated most recently by ECRI Institute on March 25, 2008. The updated information was verified by the guideline developer on August 4, 2008. This summary was updated by ECRI Institute on December 26, 2008 following the FDA advisory on Innohep (tinzaparin). This summary was updated by ECRI Institute on July 20, 2009 following the U.S. Food and Drug Administration advisory on Varenicline and Bupropion. This summary was updated by ECRI Institute on August 14, 2009 following the updated Colchicine advisory. This

summary was updated by ECRI Institute on January 5, 2010 following the U.S. Food and Drug Administration advisory on Plavix (Clopidogrel). This summary was updated by ECRI Institute on May 17, 2010 following the U.S. Food and Drug Administration advisory on Plavix (clopidogrel). This summary was updated by ECRI Institute on July 26, 2010 following the U.S. Food and Drug Administration (FDA) advisory on Proton Pump Inhibitors (PPI). This summary was updated by ECRI Institute on July 27, 2010 following the FDA drug safety communication on Heparin. This summary was updated by ECRI Institute on June 27, 2011 following the U.S. Food and Drug Administration advisory on Zocor (simvastatin). This summary was updated by ECRI Institute on April 13, 2012 following the U.S. Food and Drug Administration advisories on Statin Drugs and Statins and HIV or Hepatitis C drugs. This summary was updated by ECRI Institute on March 27, 2013. The updated information was verified by the guideline developer on May 7, 2013. This summary was updated by ECRI Institute on October 28, 2013 following the U.S. Food and Drug Administration advisory on Acetaminophen. This summary was updated by ECRI Institute on March 10, 2014 following the U.S. Food and Drug Administration advisory on Low Molecular Weight Heparins.

## Copyright Statement

This NGC summary is based on the original guideline, which is subject to the guideline developer's copyright restrictions as follows:

Copyright to the original guideline is owned by the American College of Cardiology Foundation (ACCF) and the American Heart Association, Inc. (AHA). NGC users are free to download a single copy for personal use. Reproduction without permission of the ACC/AHA guidelines is prohibited. Permissions requests should be directed to [copyright\\_permissions@acc.org](mailto:copyright_permissions@acc.org).

## Disclaimer

### NGC Disclaimer

The National Guideline Clearinghouse<sup>â„¢</sup> (NGC) does not develop, produce, approve, or endorse the guidelines represented on this site.

All guidelines summarized by NGC and hosted on our site are produced under the auspices of medical specialty societies, relevant professional associations, public or private organizations, other government agencies, health care organizations or plans, and similar entities.

Guidelines represented on the NGC Web site are submitted by guideline developers, and are screened solely to determine that they meet the [NGC Inclusion Criteria](#).

NGC, AHRQ, and its contractor ECRI Institute make no warranties concerning the content or clinical efficacy or effectiveness of the clinical practice guidelines and related materials represented on this site. Moreover, the views and opinions of developers or authors of guidelines represented on this site do not necessarily state or reflect those of NGC, AHRQ, or its contractor ECRI Institute, and inclusion or hosting of guidelines in NGC may not be used for advertising or commercial endorsement purposes.

Readers with questions regarding guideline content are directed to contact the guideline developer.